

# Linear Algebra Lecture Notes

Lecturers: Inna Capdeboscq and Damiano Testa

Warwick, January 2017

## Contents

<b>1</b>	<b>Number Systems and Fields</b>	<b>3</b>
1.1	Axioms for number systems . . . . .	3
<b>2</b>	<b>Vector Spaces</b>	<b>4</b>
2.1	Examples of vector spaces . . . . .	4
<b>3</b>	<b>Linear Independence, Spanning and Bases of Vector Spaces</b>	<b>6</b>
3.1	Linear dependence and independence . . . . .	6
3.2	Spanning vectors . . . . .	6
3.3	Bases of vector spaces . . . . .	6
3.4	Existence of a basis . . . . .	10
<b>4</b>	<b>Subspaces</b>	<b>10</b>
<b>5</b>	<b>Linear Transformations</b>	<b>13</b>
5.1	Definition and examples . . . . .	13
5.2	Kernels and images . . . . .	15
5.3	Rank and nullity . . . . .	16
5.4	Operations on linear maps . . . . .	17
<b>6</b>	<b>Matrices</b>	<b>18</b>
<b>7</b>	<b>Linear Transformations and Matrices</b>	<b>19</b>
7.1	Setting up the correspondence . . . . .	19
7.2	The correspondence between operations on linear maps and matrices . . . . .	22
7.3	Linear equations and the inverse image problem . . . . .	23
<b>8</b>	<b>Elementary Operations and the Rank of a Matrix</b>	<b>24</b>
8.1	Gauss transformations . . . . .	24
8.2	Elementary row operations . . . . .	25

8.3	Augmented Matrix . . . . .	25
8.4	Row reduction a matrix . . . . .	26
8.5	Elementary column operations . . . . .	28
8.6	The rank of a matrix . . . . .	29
8.7	Rank Criterion . . . . .	32
<b>9</b>	<b>The Inverse of a Linear Transformation and of a Matrix</b>	<b>32</b>
9.1	Definitions . . . . .	32
9.2	Matrix inversion by row reduction . . . . .	33
9.3	Elementary matrices . . . . .	35
9.4	Application to linear equations . . . . .	36
<b>10</b>	<b>The Determinant of a Matrix</b>	<b>36</b>
10.1	Definition of the determinant . . . . .	36
10.2	The effect of matrix operations on the determinant . . . . .	38
10.3	The determinant of a product . . . . .	41
10.4	Minors and cofactors . . . . .	42
10.5	The inverse of a matrix using determinants . . . . .	43
10.6	Cramer's rule for solving simultaneous equations . . . . .	44
<b>11</b>	<b>Change of Basis and Equivalent Matrices</b>	<b>44</b>
<b>12</b>	<b>Similar Matrices, Eigenvectors and Eigenvalues</b>	<b>49</b>
12.1	Similar matrices . . . . .	49
12.2	Eigenvectors and eigenvalues . . . . .	50
12.3	The scalar product - symmetric and orthogonal matrices . . . . .	55
12.4	Linear recursive sequences . . . . .	57

# 1 Number Systems and Fields

We introduce the number systems most commonly used in mathematics.

(a) The natural numbers  $\mathbb{N} = \{1, 2, 3, 4, \dots\}$ .

In  $\mathbb{N}$ , addition is possible but not subtraction; e.g.  $2 - 3 \notin \mathbb{N}$ .

(b) The integers  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, 3, \dots\}$ .

In  $\mathbb{Z}$ , addition, subtraction and multiplication are always possible, but not division; e.g.  $2/3 \notin \mathbb{Z}$ .

(c) The rational numbers  $\mathbb{Q} = \{p/q \mid p, q \in \mathbb{Z}, q \neq 0\}$ .

In  $\mathbb{Q}$ , addition, subtraction, multiplication and division (except by zero) are all possible. However,  $\sqrt{2} \notin \mathbb{Q}$ .

(d) The real numbers  $\mathbb{R}$ . These are the numbers which can be expressed as decimals. The rational numbers are those with finite or recurring decimals.

In  $\mathbb{R}$ , addition, subtraction, multiplication and division (except by zero) are still possible, and all positive numbers have square roots, but  $\sqrt{-1} \notin \mathbb{R}$ .

(e) The complex numbers  $\mathbb{C} = \{x + iy \mid x, y \in \mathbb{R}\}$ , where  $i^2 = -1$ .

In  $\mathbb{C}$ , addition, subtraction, multiplication and division (except by zero) are still possible, and all numbers have square roots. In fact all polynomial equations with coefficients in  $\mathbb{C}$  have solutions in  $\mathbb{C}$ .

## 1.1 Axioms for number systems

Laws governing the way numbers combine together are called *axioms*.

**Axioms for addition.** Let  $S$  be a number system.

A1.  $\alpha + \beta = \beta + \alpha$  for all  $\alpha, \beta \in S$ .

A2.  $(\alpha + \beta) + \gamma = \alpha + (\beta + \gamma)$  for all  $\alpha, \beta, \gamma \in S$ .

A3. There is a number  $0 \in S$  such that  $\alpha + 0 = 0 + \alpha = \alpha$  for all  $\alpha \in S$ .

A4. For each number  $\alpha \in S$  there exists a number  $-\alpha \in S$  such that  $\alpha + (-\alpha) = (-\alpha) + \alpha = 0$ .

These axioms may or may not be satisfied by a given number system  $S$ . For example, in  $\mathbb{N}$ , A1 and A2 hold but A3 and A4 do not hold. A1-A4 all hold in  $\mathbb{Z}, \mathbb{Q}, \mathbb{R}$  and  $\mathbb{C}$ .

**Axioms for multiplication.**

M1.  $\alpha.\beta = \beta.\alpha$  for all  $\alpha, \beta \in S$ .

M2.  $(\alpha.\beta).\gamma = \alpha.(\beta.\gamma)$  for all  $\alpha, \beta, \gamma \in S$ .

M3. There is a number  $1 \in S$  such that  $\alpha.1 = 1.\alpha = \alpha$  for all  $\alpha \in S$ .

M4. For each number  $\alpha \in S$  with  $\alpha \neq 0$ , there exists  $\alpha^{-1} \in S$  such that  $\alpha.\alpha^{-1} = \alpha^{-1}.\alpha = 1$ .

In  $\mathbb{N}$  and  $\mathbb{Z}$ , M1-M3 hold but M4 does not hold. M1-M4 all hold in  $\mathbb{Q}, \mathbb{R}$  and  $\mathbb{C}$ .

**Axiom relating addition and multiplication.**

D.  $(\alpha + \beta).\gamma = \alpha.\gamma + \beta.\gamma$  for all  $\alpha, \beta, \gamma \in S$ .

**Definition** A set  $S$  on which addition and multiplication are defined is called a *field*, if it satisfies each of the axioms A1, A2, A3, A4, M1, M2, M3, M4, D, and if, in addition,  $1 \neq 0$ .

Roughly speaking,  $S$  is a field if addition, subtraction, multiplication and division (except by zero) are all possible in  $S$ . We shall always use the letter  $K$  for a general field.

For example  $\mathbb{N}$  and  $\mathbb{Z}$  are not fields, but  $\mathbb{Q}, \mathbb{R}$  and  $\mathbb{C}$  are all fields.

There are many other fields, including some finite fields.

For example, for each prime number  $p$ , there is a field  $\mathbb{F}_p = \{0, 1, 2, \dots, p-1\}$  with  $p$  elements, where addition and multiplication are carried out modulo  $p$ . Thus, in  $\mathbb{F}_7$ , we have  $5 + 4 = 2$ ,  $5 \times 4 = 6$  and  $5^{-1} = 3$  because  $5 \times 3 = 1$ .

The smallest such field  $\mathbb{F}_2$  has just two elements 0 and 1, where  $1 + 1 = 0$ . This field is extremely important in Computer Science since an element of  $\mathbb{F}_2$  represents a bit of information.

Various other familiar properties of numbers, such as  $0\alpha = 0$ ,  $(-\alpha)\beta = -(\alpha\beta) = \alpha(-\beta)$ ,  $(-\alpha)(-\beta) = \alpha\beta$ ,  $(-1)\alpha = -\alpha$ , for all  $\alpha, \beta \in S$ , can be proved from the axioms.

However, occasionally you need to be careful. For example, in  $\mathbb{F}_2$  we have  $1 + 1 = 0$ , and so it is not possible to divide by 2 in this field.

## 2 Vector Spaces

**Definition** A *vector space* over a field  $K$  is a set  $V$  which has two basic operations, addition and scalar multiplication. Thus for every pair  $\mathbf{u}, \mathbf{v} \in V$ ,  $\mathbf{u} + \mathbf{v} \in V$  is defined, and for every  $\alpha \in K$ ,  $\alpha\mathbf{v} \in V$  is defined. The following axioms are satisfied for all  $\alpha, \beta \in K$  and all  $\mathbf{u}, \mathbf{v} \in V$ .

- (i) Vector addition satisfies axioms A1, A2, A3 and A4.
- (ii)  $\alpha(\mathbf{u} + \mathbf{v}) = \alpha\mathbf{u} + \alpha\mathbf{v}$ ;
- (iii)  $(\alpha + \beta)\mathbf{v} = \alpha\mathbf{v} + \beta\mathbf{v}$ ;
- (iv)  $(\alpha\beta)\mathbf{v} = \alpha(\beta\mathbf{v})$ ;
- (v)  $1\mathbf{v} = \mathbf{v}$ .

Elements of the field  $K$  will be called *scalars*. Note that we will use boldface letters like  $\mathbf{v}$  to denote vectors. The zero vector in  $V$  will be written as  $\mathbf{0}_V$ , or usually just  $\mathbf{0}$ . This is different from the zero scalar  $0 = 0_K \in K$ .

For nearly all results in this course, there is no loss in assuming that  $K$  is the field  $\mathbb{R}$  of real numbers. So you may assume this if you find it helpful to do so. Just occasionally, we will need to assume  $K = \mathbb{C}$  the field of complex numbers.

However, it is important to note that nearly all arguments in Linear Algebra use only the axioms for a field and so are valid for *any* field, which is why shall use a general field  $K$  for most of the course.

### 2.1 Examples of vector spaces

1.  $K^n = \{(\alpha_1, \alpha_2, \dots, \alpha_n) \mid \alpha_i \in K\}$ . This is the space of row vectors. Addition and scalar multiplication are defined by the obvious rules:

$$\begin{aligned}(\alpha_1, \alpha_2, \dots, \alpha_n) + (\beta_1, \beta_2, \dots, \beta_n) &= (\alpha_1 + \beta_1, \alpha_2 + \beta_2, \dots, \alpha_n + \beta_n); \\ \lambda(\alpha_1, \alpha_2, \dots, \alpha_n) &= (\lambda\alpha_1, \lambda\alpha_2, \dots, \lambda\alpha_n).\end{aligned}$$

The most familiar examples are


$$\mathbb{R}^2 = \{(x, y) \mid x, y \in \mathbb{R}\} \quad \text{and} \quad \mathbb{R}^3 = \{(x, y, z) \mid x, y, z \in \mathbb{R}\},$$

which are the points in an ordinary 2- and 3-dimensional space, equipped with a coordinate system.

Vectors in  $\mathbb{R}^2$  and  $\mathbb{R}^3$  can also be thought of as directed lines joining the origin to the points with coordinates  $(x, y)$  or  $(x, y, z)$ .



Addition of vectors is then given by the parallelogram law.



Note that  $K^1$  is essentially the same as  $K$  itself.

**2.** Let  $K[x]$  be the set of polynomials in an indeterminate  $x$  with coefficients in the field  $K$ . That is,

$$K[x] = \{\alpha_0 + \alpha_1x + \cdots + \alpha_nx^n \mid \alpha_i \in K\}.$$

Then  $K[x]$  is a vector space over  $K$ .

**3.** Let  $K[x]_{\leq n}$  be the set of polynomials over  $K$  of degree at most  $n$ , for some  $n \geq 0$ . Then  $K[x]_{\leq n}$  is also a vector space over  $K$ ; in fact it is a *subspace* of  $K[x]$ .

Note that the polynomials of degree exactly  $n$  do not form a vector space. (Why?)

**4.** Let  $K = \mathbb{R}$  and let  $V$  be the set of  $n$ -times differentiable functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  which are solutions of the differential equation

$$\lambda_0 \frac{d^n f}{dx^n} + \lambda_1 \frac{d^{n-1} f}{dx^{n-1}} + \cdots + \lambda_{n-1} \frac{df}{dx} + \lambda_n f = 0.$$

for fixed  $\lambda_0, \lambda_1, \dots, \lambda_n \in \mathbb{R}$ . Then  $V$  is a vector space over  $\mathbb{R}$ , for if  $f(x)$  and  $g(x)$  are both solutions of this equation, then so are  $f(x) + g(x)$  and  $\alpha f(x)$  for all  $\alpha \in \mathbb{R}$ .

**5.** The previous example is a space of functions. There are many such examples that are important in Analysis. For example, the set  $C^k((0, 1), \mathbb{R})$  (of all functions  $f : (0, 1) \rightarrow \mathbb{R}$  such that the  $k$ -th derivative  $f^{(k)}$  exists and is continuous) is a vector space over  $\mathbb{R}$  with the usual pointwise definitions of addition and scalar multiplication of functions.

**8.** Any  $n$  bits of information form a vector in  $\mathbb{F}_2^n$ .

Facing such a variety of vector spaces, a mathematician wants to derive useful methods of handling all these vector spaces. If we do it on a single example, say  $\mathbb{R}^8$ , how can we be certain that our methods are correct? It is only possible with the *axiomatic approach* to developing mathematics. We must use only arguments based on the vector space axioms. We have to avoid making any other assumptions. This ensures that everything we prove is valid for all vector spaces, not just the familiar ones like  $\mathbb{R}^3$ .

We shall be assuming the following additional simple properties of vectors and scalars from now on. They can all be deduced from the axioms.

- (i)  $\alpha \mathbf{0} = \mathbf{0}$  for all  $\alpha \in K$
- (ii)  $0 \mathbf{v} = \mathbf{0}$  for all  $\mathbf{v} \in V$
- (iii)  $-(\alpha \mathbf{v}) = (-\alpha) \mathbf{v} = \alpha(-\mathbf{v})$ , for all  $\alpha \in K$  and  $\mathbf{v} \in V$ .
- (iv) if  $\alpha \mathbf{v} = \mathbf{0}$  then  $\alpha = 0$  or  $\mathbf{v} = \mathbf{0}$ .

# 3 Linear Independence, Spanning and Bases of Vector Spaces

## 3.1 Linear dependence and independence

**Definition** Let  $V$  be a vector space over the field  $K$ . The vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  are said to be *linearly dependent* if there exist scalars  $\alpha_1, \alpha_2, \dots, \alpha_n \in K$ , not all zero, such that

$$\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n = \mathbf{0}.$$

$\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  are said to be *linearly independent* if they are not linearly dependent. In other words, they are linearly independent if the only scalars  $\alpha_1, \alpha_2, \dots, \alpha_n \in K$  that satisfy the above equation are  $\alpha_1 = 0, \alpha_2 = 0, \dots, \alpha_n = 0$ .

Vectors of the form  $\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n$  for  $\alpha_1, \alpha_2, \dots, \alpha_n \in K$  are called *linear combinations* of  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ .

**Examples 1.** Let  $V = \mathbb{R}^2$ ,  $\mathbf{v}_1 = (1, 3)$ ,  $\mathbf{v}_2 = (2, 5)$ .

Then  $\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 = (\alpha_1 + 2\alpha_2, 3\alpha_1 + 5\alpha_2)$ , which is equal to  $\mathbf{0} = (0, 0)$  if and only if  $\alpha_1 + 2\alpha_2 = 0$  and  $3\alpha_1 + 5\alpha_2 = 0$ . Thus we have a pair of simultaneous equations in  $\alpha_1, \alpha_2$  and the only solution is  $\alpha_1 = \alpha_2 = 0$ , so  $\mathbf{v}_1, \mathbf{v}_2$  are linearly independent.

**2.** Let  $V = \mathbb{Q}^2$ ,  $\mathbf{v}_1 = (1, 3)$ ,  $\mathbf{v}_2 = (2, 6)$ .

This time the equations are  $\alpha_1 + 2\alpha_2 = 0$  and  $3\alpha_1 + 6\alpha_2 = 0$ , and there are non-zero solutions, such as  $\alpha_1 = -2, \alpha_2 = 1$ , and so  $\mathbf{v}_1, \mathbf{v}_2$  are linearly dependent.

**Lemma 3.1**  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$  are linearly dependent if and only if either  $\mathbf{v}_1 = \mathbf{0}$  or, for some  $r$ ,  $\mathbf{v}_r$  is a linear combination of  $\mathbf{v}_1, \dots, \mathbf{v}_{r-1}$ .

**PROOF:** If  $\mathbf{v}_1 = \mathbf{0}$  then by putting  $\alpha_1 = 1$  and  $\alpha_i = 0$  for  $i > 1$  we get  $\alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n = \mathbf{0}$ , so  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$  are linearly dependent.

If  $\mathbf{v}_r$  is a linear combination of  $\mathbf{v}_1, \dots, \mathbf{v}_{r-1}$ , then  $\mathbf{v}_r = \alpha_1 \mathbf{v}_1 + \dots + \alpha_{r-1} \mathbf{v}_{r-1}$  for some  $\alpha_1, \dots, \alpha_{r-1} \in K$  and so we get  $\alpha_1 \mathbf{v}_1 + \dots + \alpha_{r-1} \mathbf{v}_{r-1} - 1 \mathbf{v}_r = \mathbf{0}$  and again  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$  are linearly dependent.

Conversely, suppose that  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$  are linearly dependent, and  $\alpha_i$  are scalars, not all zero, satisfying  $\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n = \mathbf{0}$ . Let  $r$  be maximal with  $\alpha_r \neq 0$ ; then  $\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_r \mathbf{v}_r = \mathbf{0}$ . If  $r = 1$  then  $\alpha_1 \mathbf{v}_1 = \mathbf{0}$  which, by (iv) above, is only possible if  $\mathbf{v}_1 = \mathbf{0}$ . Otherwise, we get

$$\mathbf{v}_r = -\frac{\alpha_1}{\alpha_r} \mathbf{v}_1 - \dots - \frac{\alpha_{r-1}}{\alpha_r} \mathbf{v}_{r-1}.$$

In other words,  $\mathbf{v}_r$  is a linear combination of  $\mathbf{v}_1, \dots, \mathbf{v}_{r-1}$ . □

## 3.2 Spanning vectors

**Definition** The vectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$  in  $V$  *span*  $V$  if every vector  $\mathbf{v} \in V$  is a linear combination  $\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n$  of  $\mathbf{v}_1, \dots, \mathbf{v}_n$ .

## 3.3 Bases of vector spaces

**Definition** The vectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$  in  $V$  form a *basis* of  $V$  if they are linearly independent and span  $V$ .

**Proposition 3.2** *The vectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$  form a basis of  $V$  if and only if every  $\mathbf{v} \in V$  can be written uniquely as  $\mathbf{v} = \alpha_1\mathbf{v}_1 + \alpha_2\mathbf{v}_2 + \dots + \alpha_n\mathbf{v}_n$ ; that is, the coefficients  $\alpha_1, \dots, \alpha_n$  are uniquely determined by the vector  $\mathbf{v}$ .*

PROOF: Suppose that  $\mathbf{v}_1, \dots, \mathbf{v}_n$  form a basis of  $V$ . Then they span  $V$ , so certainly every  $\mathbf{v} \in V$  can be written as  $\mathbf{v} = \alpha_1\mathbf{v}_1 + \alpha_2\mathbf{v}_2 + \dots + \alpha_n\mathbf{v}_n$ . Suppose that we also had  $\mathbf{v} = \beta_1\mathbf{v}_1 + \beta_2\mathbf{v}_2 + \dots + \beta_n\mathbf{v}_n$  for some other scalars  $\beta_i \in K$ . Then we have

$$\mathbf{0} = \mathbf{v} - \mathbf{v} = (\alpha_1 - \beta_1)\mathbf{v}_1 + (\alpha_2 - \beta_2)\mathbf{v}_2 + \dots + (\alpha_n - \beta_n)\mathbf{v}_n$$

and so

$$(\alpha_1 - \beta_1) = (\alpha_2 - \beta_2) = \dots = (\alpha_n - \beta_n) = 0$$

by linear independence of  $\mathbf{v}_1, \dots, \mathbf{v}_n$ . Hence  $\alpha_i = \beta_i$  for all  $i$ , which means that the  $\alpha_i$  are uniquely determined.

Conversely, suppose that every  $\mathbf{v} \in V$  can be written uniquely as  $\mathbf{v} = \alpha_1\mathbf{v}_1 + \alpha_2\mathbf{v}_2 + \dots + \alpha_n\mathbf{v}_n$ . Then  $\mathbf{v}_1, \dots, \mathbf{v}_n$  certainly span  $V$ . If  $\alpha_1\mathbf{v}_1 + \alpha_2\mathbf{v}_2 + \dots + \alpha_n\mathbf{v}_n = \mathbf{0}$ , then

$$\alpha_1\mathbf{v}_1 + \alpha_2\mathbf{v}_2 + \dots + \alpha_n\mathbf{v}_n = 0\mathbf{v}_1 + 0\mathbf{v}_2 + \dots + 0\mathbf{v}_n$$

and so the uniqueness assumption implies that  $\alpha_1 = \alpha_2 = \dots = \alpha_n = 0$ , and  $\mathbf{v}_1, \dots, \mathbf{v}_n$  are linearly independent. Hence they form a basis of  $V$ .  $\square$

**Examples 1.**  $(1, 0)$  and  $(0, 1)$  form a basis of  $K^2$ . This follows from Proposition 3.2, because each element  $(\alpha_1, \alpha_2) \in K^2$  can be written as  $\alpha_1(1, 0) + \alpha_2(0, 1)$ , and this expression is clearly unique.

**2.** More generally,  $(1, 0, 0)$ ,  $(0, 1, 0)$ ,  $(0, 0, 1)$  form a basis of  $K^3$ ,  $(1, 0, 0, 0)$ ,  $(0, 1, 0, 0)$ ,  $(0, 0, 1, 0)$ ,  $(0, 0, 0, 1)$  form a basis of  $K^4$  and so on. This is called the *standard basis* of  $K^n$  for  $n \in \mathbb{N}$ .

(To be precise, the standard basis of  $K^n$  is  $\mathbf{v}_1, \dots, \mathbf{v}_n$ , where  $\mathbf{v}_i$  is the vector with a 1 in the  $i$ -th position and a 0 in all other positions.)

**3.** There are many other bases of  $K^n$ . For example  $(1, 0)$ ,  $(1, 1)$  form a basis of  $K^2$ , because  $(\alpha_1, \alpha_2) = (\alpha_1 - \alpha_2)(1, 0) + \alpha_2(1, 1)$ , and this expression is unique. In fact, any two non-zero vectors such that one is not a scalar multiple of the other form a basis for  $K^2$ .

**4.** Not every vector space has a finite basis. For example, let  $K[x]$  be the space of polynomials in  $x$  with coefficients in  $K$ . Let  $p_1(x), p_2(x), \dots, p_n(x)$  be any finite collection of polynomials in  $K[x]$ . Then if  $d$  is the maximum degree of  $p_1(x), p_2(x), \dots, p_n(x)$ , any linear combination of  $p_1(x), p_2(x), \dots, p_n(x)$  has degree at most  $d$ , and so  $p_1(x), p_2(x), \dots, p_n(x)$  cannot span  $K[x]$ . It is easy to see that the infinite sequence of vectors  $1, x, x^2, x^3, \dots, x^n, \dots$  is a basis of  $K[x]$ .

A vector space with a finite basis is called *finite-dimensional*. In fact, nearly all of this course will be about finite-dimensional spaces, but it is important to remember that these are not the only examples. The spaces of functions mentioned in Example 5. of Section 2 typically have uncountably infinite dimension.

**Theorem 3.3** (The basis theorem) *Suppose that  $\mathbf{v}_1, \dots, \mathbf{v}_m$  and  $\mathbf{w}_1, \dots, \mathbf{w}_n$  are both finite bases of the vector space  $V$ . Then  $m = n$ . In other words, all finite bases of  $V$  contain the same number of vectors.*

The proof of this theorem is quite tricky and uses the concept of *sifting* which we introduce after the next lemma.

**Definition** The number  $n$  of vectors in a basis of the finite-dimensional vector space  $V$  is called the *dimension* of  $V$  and we write  $\dim(V) = n$ .

Thus, as we might expect,  $K^n$  has dimension  $n$ .  $K[x]$  is infinite-dimensional, but the space  $K[x]_{\leq n}$  of polynomials of degree at most  $n$  has basis  $1, x, x^2, \dots, x^n$ , so its dimension is  $n + 1$  (not  $n$ ).

Note that the dimension of  $V$  depends on the field  $K$ . Thus the complex numbers  $\mathbb{C}$  can be considered as

- a space of dimension 1 over  $\mathbb{C}$ ,
- a space of dimension 2 over  $\mathbb{R}$ , where  $1, i$  is a basis for  $\mathbb{C}$  over  $\mathbb{R}$ ,
- a space of infinite dimension over  $\mathbb{Q}$ .

**Lemma 3.4** *Suppose that the vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n, \mathbf{w}$  span  $V$  and that  $\mathbf{w}$  is a linear combination of  $\mathbf{v}_1, \dots, \mathbf{v}_n$ . Then  $\mathbf{v}_1, \dots, \mathbf{v}_n$  span  $V$ .*

PROOF: Since  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n, \mathbf{w}$  span  $V$ , any vector  $\mathbf{v} \in V$  can be written as

$$\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n + \beta \mathbf{w},$$

But  $\mathbf{w}$  is a linear combination of  $\mathbf{v}_1, \dots, \mathbf{v}_n$ , so  $\mathbf{w} = \gamma_1 \mathbf{v}_1 + \dots + \gamma_n \mathbf{v}_n$  for some scalars  $\gamma_i$ , and hence

$$\mathbf{v} = (\alpha_1 + \beta \gamma_1) \mathbf{v}_1 + \dots + (\alpha_n + \beta \gamma_n) \mathbf{v}_n$$

is a linear combination of  $\mathbf{v}_1, \dots, \mathbf{v}_n$ , which therefore span  $V$ . □

There is an important process, which we shall call *sifting*, which can be applied to any sequence of vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  in a vector space  $V$ . We consider each vector  $\mathbf{v}_i$  in turn. If it is zero, or a linear combination of the preceding vectors  $\mathbf{v}_1, \dots, \mathbf{v}_{i-1}$ , then we remove it from the list.

**Example** Let us sift the following sequence of vectors in  $\mathbb{R}^3$ .

$$\begin{array}{llll} \mathbf{v}_1 = (0, 0, 0) & \mathbf{v}_2 = (1, 1, 1) & \mathbf{v}_3 = (2, 2, 2) & \mathbf{v}_4 = (1, 0, 0) \\ \mathbf{v}_5 = (3, 2, 2) & \mathbf{v}_6 = (0, 0, 0) & \mathbf{v}_7 = (1, 1, 0) & \mathbf{v}_8 = (0, 0, 1) \end{array}$$

$\mathbf{v}_1 = \mathbf{0}$ , so we remove it.  $\mathbf{v}_2$  is non-zero so it stays.  $\mathbf{v}_3 = 2\mathbf{v}_2$  so it is removed.  $\mathbf{v}_4$  is clearly not a linear combination of  $\mathbf{v}_2$ , so it stays.

We have to decide next whether  $\mathbf{v}_5$  is a linear combination of  $\mathbf{v}_2, \mathbf{v}_4$ . If so, then  $(3, 2, 2) = \alpha_1(1, 1, 1) + \alpha_2(1, 0, 0)$ , which (fairly obviously) has the solution  $\alpha_1 = 2, \alpha_2 = 1$ , so remove  $\mathbf{v}_5$ . Then  $\mathbf{v}_6 = \mathbf{0}$  so that is removed too.

Next we try  $\mathbf{v}_7 = (1, 1, 0) = \alpha_1(1, 1, 1) + \alpha_2(1, 0, 0)$ , and looking at the three components, this reduces to the three equations

$$1 = \alpha_1 + \alpha_2; \quad 1 = \alpha_1; \quad 0 = \alpha_1.$$

The second and third of these equations contradict each other, and so there is no solution. Hence  $\mathbf{v}_7$  is not a linear combination of  $\mathbf{v}_2, \mathbf{v}_4$ , and it stays.

Finally, we need to try

$$\mathbf{v}_8 = (0, 0, 1) = \alpha_1(1, 1, 1) + \alpha_2(1, 0, 0) + \alpha_3(1, 1, 0)$$

leading to the three equations

$$0 = \alpha_1 + \alpha_2 + \alpha_3 \quad 0 = \alpha_1 + \alpha_3; \quad 1 = \alpha_1$$

and solving these in the normal way, we find a solution  $\alpha_1 = 1, \alpha_2 = 0, \alpha_3 = -1$ . Thus we delete  $\mathbf{v}_8$  and we are left with just  $\mathbf{v}_2, \mathbf{v}_4, \mathbf{v}_7$ .

Of course, the vectors that are removed during the sifting process depends very much on the order of the list of vectors. For example, if  $\mathbf{v}_8$  had come at the beginning of the list rather than at the end, then we would have kept it.

**Theorem 3.5** *Suppose that the vectors  $\mathbf{v}_1, \dots, \mathbf{v}_r$  span the vector space  $V$ . Then there is a subsequence of  $\mathbf{v}_1, \dots, \mathbf{v}_r$  which forms a basis of  $V$ .*

PROOF: We sift the vectors  $\mathbf{v}_1, \dots, \mathbf{v}_r$ . The vectors that we remove are linear combinations of the preceding vectors, and so by Lemma 3.4, the remaining vectors still span  $V$ . After sifting, no vector is zero or a linear combination of the preceding vectors (or it would have been removed), so by Lemma 3.1, the remaining vectors are linearly independent. Hence they form a basis of  $V$ .  $\square$

The theorem tells us that any vector space with a finite spanning set is finite-dimensional, indeed the spanning set contains a basis. We now prove the dual result: any linearly independent set is contained in a basis.

**Theorem 3.6** *Let  $V$  be a vector space over  $K$  which has a finite spanning set, and suppose that the vectors  $\mathbf{v}_1, \dots, \mathbf{v}_r$  are linearly independent in  $V$ . Then we can extend the sequence to a basis  $\mathbf{v}_1, \dots, \mathbf{v}_n$  of  $V$ , where  $n \geq r$ .*

PROOF: Suppose that  $\mathbf{w}_1, \dots, \mathbf{w}_q$  is a spanning set for  $V$ . We sift the combined sequence

$$\mathbf{v}_1, \dots, \mathbf{v}_r, \mathbf{w}_1, \dots, \mathbf{w}_q.$$

Since  $\mathbf{w}_1, \dots, \mathbf{w}_q$  span  $V$ , the whole sequence spans  $V$ . Sifting results in a basis for  $V$  as in the proof of Theorem 3.5. Since  $\mathbf{v}_1, \dots, \mathbf{v}_r$  are linearly independent, none of them can be a linear combination of the preceding vectors, and hence none of the  $\mathbf{v}_i$  are deleted in the sifting process. Thus the resulting basis contains  $\mathbf{v}_1, \dots, \mathbf{v}_r$ .  $\square$

**Example** The vectors  $\mathbf{v}_1 = (1, 2, 0, 2), \mathbf{v}_2 = (0, 1, 0, 2)$  are linearly independent in  $\mathbb{R}^4$ . Let us extend them to a basis of  $\mathbb{R}^4$ . The easiest thing is to append the standard basis of  $\mathbb{R}^4$ , giving the combined list of vectors

$$\begin{array}{lll} \mathbf{v}_1 = (1, 2, 0, 2), & \mathbf{v}_2 = (0, 1, 0, 2), & \mathbf{w}_1 = (1, 0, 0, 0), \\ \mathbf{w}_2 = (0, 1, 0, 0), & \mathbf{w}_3 = (0, 0, 1, 0), & \mathbf{w}_4 = (0, 0, 0, 1), \end{array}$$

which we shall sift. We find that  $(1, 0, 0, 0) = \alpha_1(1, 2, 0, 2) + \alpha_2(0, 1, 0, 2)$  has no solution, so  $\mathbf{w}_1$  stays. However,  $\mathbf{w}_2 = \mathbf{v}_1 - \mathbf{v}_2 - \mathbf{w}_1$  so  $\mathbf{w}_2$  is deleted. It is clear that  $\mathbf{w}_3$  is not a linear combination of  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{w}_1$ , because all of those have a 0 in their third component. Hence  $\mathbf{w}_3$  remains. Now we have four linearly independent vectors, so must have a basis at this stage, and we can stop the sifting early. The resulting basis is

$$\mathbf{v}_1 = (1, 2, 0, 2), \quad \mathbf{v}_2 = (0, 1, 0, 2), \quad \mathbf{w}_1 = (1, 0, 0, 0), \quad \mathbf{w}_3 = (0, 0, 1, 0).$$

We are now ready to prove Theorem 3.3. Since bases of  $V$  are both linearly independent and span  $V$ , the following proposition implies that any two bases contain the same number of vectors.

**Proposition 3.7** (The interchange lemma) *Suppose that vectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$  span  $V$  and that vectors  $\mathbf{w}_1, \dots, \mathbf{w}_m \in V$  are linearly independent. Then  $m \leq n$ .*

PROOF: The idea is to place the  $\mathbf{w}_i$  one by one in front of the sequence  $\mathbf{v}_1, \dots, \mathbf{v}_n$ , sifting each time.

Since  $\mathbf{v}_1, \dots, \mathbf{v}_n$  span  $V$ ,  $\mathbf{w}_1, \mathbf{v}_1, \dots, \mathbf{v}_n$  are linearly dependent, so when we sift, at least one  $\mathbf{v}_j$  is deleted. We then place  $\mathbf{w}_2$  in front of the resulting sequence and sift again. Then we put  $\mathbf{w}_3$  in front of the result, and sift again, and carry on doing this for each  $\mathbf{w}_i$  in turn. Since  $\mathbf{w}_1, \dots, \mathbf{w}_m$  are linearly independent none of them are ever deleted. Each time we place a vector in front of a sequence which spans  $V$ , and so the extended sequence is linearly dependent, and hence at least one  $\mathbf{v}_j$  gets eliminated each time.

But in total, we append  $m$  vectors  $\mathbf{w}_i$ , and each time at least one  $\mathbf{v}_j$  is eliminated, so we must have  $m \leq n$ .  $\square$

**Corollary 3.8** *Let  $V$  be a vector space of dimension  $n$  over  $K$ . Then any  $n$  vectors which span  $V$  form a basis of  $V$ , and no  $n - 1$  vectors can span  $V$ .*

PROOF: After sifting a spanning sequence as in the proof of Theorem 3.5, the remaining vectors form a basis, so by Theorem 3.3, there must be precisely  $n = \dim(V)$  vectors remaining. The result is now clear.  $\square$

**Corollary 3.9** *Let  $V$  be a vector space of dimension  $n$  over  $K$ . Then any  $n$  linearly independent vectors form a basis of  $V$  and no  $n + 1$  vectors can be linearly independent.*

PROOF: By Theorem 3.6 any linearly independent set is contained in a basis but by Theorem 3.3, there must be precisely  $n = \dim(V)$  vectors in the extended set. The result is now clear.  $\square$

(**Remark for the observant reader:** Notice that this Corollary shows that a vector space  $V$  cannot have both a finite and an infinite basis.)

### 3.4 Existence of a basis

Although we have studied bases quite carefully in the previous section, we have not addressed the following fundamental question. Let  $V$  be a vector space. Does it contain a basis?

Theorem 3.5 gives a partial answer that is good for many practical purposes. Let us formulate it as a corollary.

**Corollary 3.10** *If a non-trivial vector space  $V$  is spanned by a finite number of vectors, then it has a basis.*

In fact, the following general theorem is true.

**Theorem 3.11** *Any vector space  $V$  has a basis.*

Theorem 3.11 will not be proved in this course, thus, not assessed on the exam. Its proof requires Zorn's lemma that would lead us too deep into axiomatic Set Theory.

While Theorem 3.5 gives an algorithm for constructing bases, Theorem 3.11 is merely an existence theorem. For instance, the vector space of all sequences  $V = \{(a_1, a_2, \dots) \mid a_i \in \mathbb{R}\}$  has a basis by Theorem 3.11 but no such basis is explicitly known.

## 4 Subspaces

Let  $V$  be a vector space over the field  $K$ .

**Definition** A subspace of  $V$  is a non-empty subset  $W \subseteq V$  such that

$$\mathbf{u}, \mathbf{v} \in W \Rightarrow \mathbf{u} + \mathbf{v} \in W \quad \text{and} \quad \mathbf{v} \in W, \alpha \in K \Rightarrow \alpha \mathbf{v} \in W.$$

These two conditions can be replaced with a single condition

$$\mathbf{u}, \mathbf{v} \in W, \alpha, \beta \in K \Rightarrow \alpha \mathbf{u} + \beta \mathbf{v} \in W.$$

A subspace  $W$  is itself a vector space over  $K$  under the operations of vector addition and scalar multiplication in  $V$ . Notice that all vector space axioms of  $W$  hold automatically. (They are inherited from  $V$ .)

For any vector space  $V$ ,  $V$  is always a subspace of itself. Subspaces other than  $V$  are sometimes called *proper* subspaces. We also always have a subspace  $\{\mathbf{0}\}$  consisting of the zero vector alone. This is called the *trivial* subspace, and its dimension is 0, because it has no linearly independent sets of vectors at all.

**Proposition 4.1** *If  $W_1$  and  $W_2$  are subspaces of  $V$  then so is  $W_1 \cap W_2$ .*

PROOF: Let  $\mathbf{u}, \mathbf{v} \in W_1 \cap W_2$  and  $\alpha \in K$ . Then  $\mathbf{u} + \mathbf{v} \in W_1$  (because  $W_1$  is a subspace) and  $\mathbf{u} + \mathbf{v} \in W_2$  (because  $W_2$  is a subspace). Hence  $\mathbf{u} + \mathbf{v} \in W_1 \cap W_2$ . Similarly, we get  $\alpha \mathbf{v} \in W_1 \cap W_2$ , so  $W_1 \cap W_2$  is a subspace of  $V$ .  $\square$

**Warning!** It is **not** necessarily true that  $W_1 \cup W_2$  is a subspace.

**Example** Let  $V = \mathbb{R}^2$ , let  $W_1 = \{(\alpha, 0) \mid \alpha \in \mathbb{R}\}$  and  $W_2 = \{(0, \alpha) \mid \alpha \in \mathbb{R}\}$ . Then  $W_1, W_2$  are subspaces of  $V$ , but  $W_1 \cup W_2$  is not a subspace, because  $(1, 0), (0, 1) \in W_1 \cup W_2$ , but  $(1, 0) + (0, 1) = (1, 1) \notin W_1 \cup W_2$ .

Note that any subspace of  $V$  that contains  $W_1$  and  $W_2$  has to contain all vectors of the form  $\mathbf{u} + \mathbf{v}$  for  $\mathbf{u} \in W_1, \mathbf{v} \in W_2$ .

**Definition** Let  $W_1, W_2$  be subspaces of the vector space  $V$ . Then  $W_1 + W_2$  is defined to be the set of vectors  $\mathbf{v} \in V$  such that  $\mathbf{v} = \mathbf{w}_1 + \mathbf{w}_2$  for some  $\mathbf{w}_1 \in W_1, \mathbf{w}_2 \in W_2$ . Or, if you prefer,  $W_1 + W_2 = \{\mathbf{w}_1 + \mathbf{w}_2 \mid \mathbf{w}_1 \in W_1, \mathbf{w}_2 \in W_2\}$ .

Do not confuse  $W_1 + W_2$  with  $W_1 \cup W_2$ .

**Proposition 4.2** *If  $W_1, W_2$  are subspaces of  $V$  then so is  $W_1 + W_2$ . In fact, it is the smallest subspace that contains both  $W_1$  and  $W_2$ .*

PROOF: Let  $\mathbf{u}, \mathbf{v} \in W_1 + W_2$ . Then  $\mathbf{u} = \mathbf{u}_1 + \mathbf{u}_2$  for some  $\mathbf{u}_1 \in W_1, \mathbf{u}_2 \in W_2$  and  $\mathbf{v} = \mathbf{v}_1 + \mathbf{v}_2$  for some  $\mathbf{v}_1 \in W_1, \mathbf{v}_2 \in W_2$ . Then  $\mathbf{u} + \mathbf{v} = (\mathbf{u}_1 + \mathbf{v}_1) + (\mathbf{u}_2 + \mathbf{v}_2) \in W_1 + W_2$ . Similarly, if  $\alpha \in K$  then  $\alpha \mathbf{v} = \alpha \mathbf{v}_1 + \alpha \mathbf{v}_2 \in W_1 + W_2$ . Thus  $W_1 + W_2$  is a subspace of  $V$ .

Any subspace of  $V$  that contains both  $W_1$  and  $W_2$  must contain  $W_1 + W_2$ , so it is the smallest such subspace.  $\square$

**Theorem 4.3** *Let  $V$  be a finite-dimensional vector space, and let  $W_1, W_2$  be subspaces of  $V$ . Then*

$$\dim(W_1 + W_2) = \dim(W_1) + \dim(W_2) - \dim(W_1 \cap W_2).$$

PROOF: First note that any subspace  $W$  of  $V$  is finite-dimensional. This follows from Corollary 3.9, because a largest linearly independent subset of  $W$  contains at most  $\dim(V)$  vectors, and such a subset must be a basis of  $W$ .

Let  $\dim(W_1 \cap W_2) = r$  and let  $\mathbf{e}_1, \dots, \mathbf{e}_r$  be a basis of  $W_1 \cap W_2$ . Then  $\mathbf{e}_1, \dots, \mathbf{e}_r$  is a linearly independent set of vectors, so by Theorem 3.6 it can be extended to a basis  $\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{f}_1, \dots, \mathbf{f}_s$

of  $W_1$  where  $\dim(W_1) = r + s$ , and it can also be extended to a basis  $\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{g}_1, \dots, \mathbf{g}_t$  of  $W_2$ , where  $\dim(W_2) = r + t$ .

To prove the theorem, we need to show that  $\dim(W_1 + W_2) = r + s + t$ , and to do this, we shall show that

$$\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{f}_1, \dots, \mathbf{f}_s, \mathbf{g}_1, \dots, \mathbf{g}_t$$

is a basis of  $W_1 + W_2$ . Certainly they all lie in  $W_1 + W_2$ .

First we show that they span  $W_1 + W_2$ . Any  $\mathbf{v} \in W_1 + W_2$  is equal to  $\mathbf{w}_1 + \mathbf{w}_2$  for some  $\mathbf{w}_1 \in W_1$ ,  $\mathbf{w}_2 \in W_2$ . So we can write

$$\mathbf{w}_1 = \alpha_1 \mathbf{e}_1 + \dots + \alpha_r \mathbf{e}_r + \beta_1 \mathbf{f}_1 + \dots + \beta_s \mathbf{f}_s$$

for some scalars  $\alpha_i, \beta_j \in K$ , and

$$\mathbf{w}_2 = \gamma_1 \mathbf{e}_1 + \dots + \gamma_r \mathbf{e}_r + \delta_1 \mathbf{g}_1 + \dots + \delta_t \mathbf{g}_t$$

for some scalars  $\gamma_i, \delta_j \in K$ . Then

$$\mathbf{v} = (\alpha_1 + \gamma_1) \mathbf{e}_1 + \dots + (\alpha_r + \gamma_r) \mathbf{e}_r + \beta_1 \mathbf{f}_1 + \dots + \beta_s \mathbf{f}_s + \delta_1 \mathbf{g}_1 + \dots + \delta_t \mathbf{g}_t$$

and so  $\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{f}_1, \dots, \mathbf{f}_s, \mathbf{g}_1, \dots, \mathbf{g}_t$  span  $W_1 + W_2$ .

Finally we have to show that  $\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{f}_1, \dots, \mathbf{f}_s, \mathbf{g}_1, \dots, \mathbf{g}_t$  are linearly independent. Suppose that

$$\alpha_1 \mathbf{e}_1 + \dots + \alpha_r \mathbf{e}_r + \beta_1 \mathbf{f}_1 + \dots + \beta_s \mathbf{f}_s + \delta_1 \mathbf{g}_1 + \dots + \delta_t \mathbf{g}_t = \mathbf{0}$$

for some scalars  $\alpha_i, \beta_j, \delta_k \in K$ . Then

$$\alpha_1 \mathbf{e}_1 + \dots + \alpha_r \mathbf{e}_r + \beta_1 \mathbf{f}_1 + \dots + \beta_s \mathbf{f}_s = -\delta_1 \mathbf{g}_1 - \dots - \delta_t \mathbf{g}_t \quad (*)$$

The left-hand-side of this equation lies in  $W_1$  and the right-hand-side of this equation lies in  $W_2$ . Since the two sides are equal, both must in fact lie in  $W_1 \cap W_2$ . Since  $\mathbf{e}_1, \dots, \mathbf{e}_r$  is a basis of  $W_1 \cap W_2$ , we can write

$$-\delta_1 \mathbf{g}_1 - \dots - \delta_t \mathbf{g}_t = \gamma_1 \mathbf{e}_1 + \dots + \gamma_r \mathbf{e}_r$$

for some  $\gamma_i \in K$ , and so

$$\gamma_1 \mathbf{e}_1 + \dots + \gamma_r \mathbf{e}_r + \delta_1 \mathbf{g}_1 + \dots + \delta_t \mathbf{g}_t = \mathbf{0}.$$

But,  $\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{g}_1, \dots, \mathbf{g}_t$  form a basis of  $W_2$ , so they are linearly independent, and hence  $\gamma_i = 0$  for  $1 \leq i \leq r$  and  $\delta_i = 0$  for  $1 \leq i \leq t$ . But now, from the equation (\*) above, we get

$$\alpha_1 \mathbf{e}_1 + \dots + \alpha_r \mathbf{e}_r + \beta_1 \mathbf{f}_1 + \dots + \beta_s \mathbf{f}_s = \mathbf{0}.$$

Now  $\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{f}_1, \dots, \mathbf{f}_s$  form a basis of  $W_1$ , so they are linearly independent, and hence  $\alpha_i = 0$  for  $1 \leq i \leq r$  and  $\beta_i = 0$  for  $1 \leq i \leq s$ . Thus  $\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{f}_1, \dots, \mathbf{f}_s, \mathbf{g}_1, \dots, \mathbf{g}_t$  are linearly independent, which completes the proof that they form a basis of  $W_1 + W_2$ .

Hence

$$\dim(W_1 + W_2) = r + s + t = (r + s) + (r + t) - r = \dim(W_1) + \dim(W_2) - \dim(W_1 \cap W_2).$$

□

**Proposition 4.4** *Let  $\mathbf{v}_1, \dots, \mathbf{v}_n$  be vectors in the vector space  $V$ . Then the set of all linear combinations  $\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n$  of  $\mathbf{v}_1, \dots, \mathbf{v}_n$  forms a subspace of  $V$ .*

The proof of this is completely routine and will be omitted. The subspace in this proposition is known as the subspace *spanned* by  $\mathbf{v}_1, \dots, \mathbf{v}_n$ .

**Definition** Two subspaces  $W_1, W_2$  of  $V$  are called *complementary* if  $W_1 \cap W_2 = \{\mathbf{0}\}$  and  $W_1 + W_2 = V$ .

**Proposition 4.5** *Let  $W_1, W_2$  be subspaces of  $V$ . Then  $W_1, W_2$  are complementary subspaces if and only if each vector in  $\mathbf{v} \in V$  can be written uniquely as  $\mathbf{v} = \mathbf{w}_1 + \mathbf{w}_2$  with  $\mathbf{w}_1 \in W_1$  and  $\mathbf{w}_2 \in W_2$ .*

PROOF: Suppose first that  $W_1, W_2$  are complementary subspaces and let  $\mathbf{v} \in V$ . Then  $W_1 + W_2 = V$ , so we can find  $\mathbf{w}_1 \in W_1$  and  $\mathbf{w}_2 \in W_2$  with  $\mathbf{v} = \mathbf{w}_1 + \mathbf{w}_2$ . If we also had  $\mathbf{v} = \mathbf{w}'_1 + \mathbf{w}'_2$  with  $\mathbf{w}'_1 \in W_1$ ,  $\mathbf{w}'_2 \in W_2$ , then we would have  $\mathbf{w}_1 - \mathbf{w}'_1 = \mathbf{w}'_2 - \mathbf{w}_2$ . The left-hand-side lies in  $W_1$  and the right-hand-side lies in  $W_2$ , and so both sides (being equal) must lie in  $W_1 \cap W_2 = \{\mathbf{0}\}$ . Hence both sides are zero, which means  $\mathbf{w}_1 = \mathbf{w}'_1$  and  $\mathbf{w}_2 = \mathbf{w}'_2$ , so the expression is unique.

Conversely, suppose that every  $\mathbf{v} \in V$  can be written uniquely as  $\mathbf{v} = \mathbf{w}_1 + \mathbf{w}_2$  with  $\mathbf{w}_1 \in W_1$  and  $\mathbf{w}_2 \in W_2$ . Then certainly  $W_1 + W_2 = V$ . If  $\mathbf{v}$  was a non-zero vector in  $W_1 \cap W_2$ , then in fact  $\mathbf{v}$  would have two distinct expressions as  $\mathbf{w}_1 + \mathbf{w}_2$  with  $\mathbf{w}_1 \in W_1$  and  $\mathbf{w}_2 \in W_2$ , one with  $\mathbf{w}_1 = \mathbf{v}$ ,  $\mathbf{w}_2 = \mathbf{0}$  and the other with  $\mathbf{w}_1 = \mathbf{0}$ ,  $\mathbf{w}_2 = \mathbf{v}$ . Hence  $W_1 \cap W_2 = \{\mathbf{0}\}$ , and  $W_1$  and  $W_2$  are complementary.  $\square$

**Examples 1.** As above, let  $V = \mathbb{R}^2$ ,  $W_1 = \{(\alpha, 0) \mid \alpha \in \mathbb{R}\}$  and  $W_2 = \{(0, \alpha) \mid \alpha \in \mathbb{R}\}$ . Then  $W_1$  and  $W_2$  are complementary subspaces.

2. As above, let  $V = \mathbb{R}^3$ ,  $W_1 = \{(\alpha, 0, 0) \mid \alpha \in \mathbb{R}\}$  and  $W_2 = \{(0, \alpha, \beta) \mid \alpha, \beta \in \mathbb{R}\}$ . Then  $W_1$  and  $W_2$  are complementary subspaces.

There are many other examples.

3. Let  $V = \mathbb{R}^2$ ,  $W_1 = \{(\alpha, \alpha) \mid \alpha \in \mathbb{R}\}$  and  $W_2 = \{(-\alpha, \alpha) \mid \alpha \in \mathbb{R}\}$ . Then  $W_1$  and  $W_2$  are complementary subspaces.

## 5 Linear Transformations

When you study sets the notion of function is extremely important. There is little to say about a single isolated set while functions allow you to link different sets. Similarly in Linear Algebra, a single isolated vector space is not the end of the story. We have to connect different vector spaces by functions. However, a function having little regard to the vector space operations may be of little value.

### 5.1 Definition and examples

**Definition** Let  $U, V$  be two vector spaces over the same field  $K$ . A *linear transformation* or *linear map*  $T$  from  $U$  to  $V$  is a function  $T : U \rightarrow V$  such that

- (i)  $T(\mathbf{u}_1 + \mathbf{u}_2) = T(\mathbf{u}_1) + T(\mathbf{u}_2)$  for all  $\mathbf{u}_1, \mathbf{u}_2 \in U$ ;
- (ii)  $T(\alpha \mathbf{u}) = \alpha T(\mathbf{u})$  for all  $\alpha \in K$  and  $\mathbf{u} \in U$ .

We shall usually call these linear maps (because this is shorter), although linear transformation is the standard name. Notice that the two conditions for linearity are equivalent to a single condition

$$T(\alpha \mathbf{u}_1 + \beta \mathbf{u}_2) = \alpha T(\mathbf{u}_1) + \beta T(\mathbf{u}_2) \text{ for all } \mathbf{u}_1, \mathbf{u}_2 \in U, \alpha, \beta \in K.$$

First a couple of trivial consequences of the definition:

**Lemma 5.1** Let  $T : U \rightarrow V$  be a linear map. Then


- (i)  $T(\mathbf{0}_U) = \mathbf{0}_V$ ;
- (ii)  $T(-\mathbf{u}) = -T(\mathbf{u})$  for all  $\mathbf{u} \in U$ .

PROOF: (i)  $T(\mathbf{0}_U) = T(\mathbf{0}_U + \mathbf{0}_U) = T(\mathbf{0}_U) + T(\mathbf{0}_U)$ , so  $T(\mathbf{0}_U) = \mathbf{0}_V$ .

(ii) Just put  $\alpha = -1$  in the definition of linear map. □


**Examples** Many familiar geometrical transformations, such as projections, rotations, reflections and magnifications are linear maps, and the first three examples below are of this kind. Note, however, that a nontrivial translation is not a linear map, because it does not satisfy  $T(\mathbf{0}_U) = \mathbf{0}_V$ .

1. Let  $U = \mathbb{R}^3$ ,  $V = \mathbb{R}^2$  and define  $T : U \rightarrow V$  by  $T((\alpha, \beta, \gamma)) = (\alpha, \beta)$ . Then  $T$  is a linear map. This type of map is known as a *projection*, because of the geometrical interpretation.



(**Note:** In future we shall just write  $T(\alpha, \beta, \gamma)$  instead of  $T((\alpha, \beta, \gamma))$ .)

2. Let  $U = V = \mathbb{R}^2$ . We interpret  $\mathbf{v}$  in  $\mathbb{R}^2$  as a directed line vector from  $\mathbf{0}$  to  $\mathbf{v}$  (see the examples in Section 2), and let  $T(\mathbf{v})$  be the vector obtained by rotating  $\mathbf{v}$  through an angle  $\theta$  anti-clockwise about the origin.




It is easy to see geometrically that  $T(\mathbf{u}_1 + \mathbf{u}_2) = T(\mathbf{u}_1) + T(\mathbf{u}_2)$  and  $T(\alpha\mathbf{u}) = \alpha T(\mathbf{u})$  (because everything is simply rotated about the origin), and so  $T$  is a linear map. By considering the unit vectors, we have  $T(1, 0) = (\cos \theta, \sin \theta)$  and  $T(0, 1) = (-\sin \theta, \cos \theta)$ , and hence

$$T(\alpha, \beta) = \alpha T(1, 0) + \beta T(0, 1) = (\alpha \cos \theta - \beta \sin \theta, \alpha \sin \theta + \beta \cos \theta).$$

(*Exercise:* Show this directly.)

3. Let  $U = V = \mathbb{R}^2$  again. Now let  $T(\mathbf{v})$  be the vector resulting from reflecting  $\mathbf{v}$  through a line through the origin that makes an angle  $\theta/2$  with the  $x$ -axis.



This is again a linear map. We find that  $T(1, 0) = (\cos \theta, \sin \theta)$  and  $T(0, 1) = (\sin \theta, -\cos \theta)$ , and so

$$T(\alpha, \beta) = \alpha T(1, 0) + \beta T(0, 1) = (\alpha \cos \theta + \beta \sin \theta, \alpha \sin \theta - \beta \cos \theta).$$

4. Let  $U = V = \mathbb{R}[x]$ , the set of polynomials over  $\mathbb{R}$ , and let  $T$  be differentiation; i.e.  $T(p(x)) = p'(x)$  for  $p \in \mathbb{R}[x]$ . This is easily seen to be a linear map.

5. Let  $U = K[x]$ , the set of polynomials over  $K$ . Every  $\alpha \in K$  gives rise to two linear maps, shift  $S_\alpha : U \rightarrow U$ ,  $S_\alpha(f(x)) = f(x - \alpha)$  and evaluation  $E_\alpha : U \rightarrow K$ ,  $E_\alpha(f(x)) = f(\alpha)$ .

The next two examples seem dull but are important!

6. For any vector space  $V$ , we define the identity map  $I_V : V \rightarrow V$  by  $I_V(\mathbf{v}) = \mathbf{v}$  for all  $\mathbf{v} \in V$ . This is a linear map.

7. For any vector spaces  $U, V$  over the field  $K$ , we define the zero map  $\mathbf{0}_{U,V} : U \rightarrow V$  by  $\mathbf{0}_{U,V}(\mathbf{u}) = \mathbf{0}_V$  for all  $\mathbf{u} \in U$ . This is also a linear map.

**Proposition 5.2** (Linear maps are uniquely determined by their action on a basis.) *Let  $U, V$  be vector spaces over  $K$ , let  $\mathbf{u}_1, \dots, \mathbf{u}_n$  be a basis of  $U$  and let  $\mathbf{v}_1, \dots, \mathbf{v}_n$  be any sequence of  $n$  vectors in  $V$ . Then there is a unique linear map  $T : U \rightarrow V$  with  $T(\mathbf{u}_i) = \mathbf{v}_i$  for  $1 \leq i \leq n$ .*

PROOF: Let  $\mathbf{u} \in U$ . Then, since  $\mathbf{u}_1, \dots, \mathbf{u}_n$  is a basis of  $U$ , by Proposition 3.2, there exist uniquely determined  $\alpha_1, \dots, \alpha_n \in K$  with  $\mathbf{u} = \alpha_1 \mathbf{u}_1 + \dots + \alpha_n \mathbf{u}_n$ . Hence, if  $T$  exists at all, then we must have

$$T(\mathbf{u}) = T(\alpha_1 \mathbf{u}_1 + \dots + \alpha_n \mathbf{u}_n) = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n,$$

and so  $T$  is uniquely determined.

On the other hand, it is routine to check that the map  $T : U \rightarrow V$  defined by the above equation is indeed a linear map, so  $T$  does exist and is unique.  $\square$

## 5.2 Kernels and images

**Definition** Let  $T : U \rightarrow V$  be a linear map. The *image* of  $T$ , written as  $\text{im}(T)$  is defined to be the set of vectors  $\mathbf{v} \in V$  such that  $\mathbf{v} = T(\mathbf{u})$  for some  $\mathbf{u} \in U$ .

The *kernel* of  $T$ , written as  $\text{ker}(T)$  is defined to be the set of vectors  $\mathbf{u} \in U$  such that  $T(\mathbf{u}) = \mathbf{0}_V$ . Or, if you prefer:

$$\text{im}(T) = \{T(\mathbf{u}) \mid \mathbf{u} \in U\}; \quad \text{ker}(T) = \{\mathbf{u} \in U \mid T(\mathbf{u}) = \mathbf{0}_V\}.$$

**Examples** Let us consider Examples 1. – 6. above.

In 1.,  $\text{ker}(T) = \{(0, 0, \gamma) \mid \gamma \in \mathbb{R}\}$ , and  $\text{im}(T) = \mathbb{R}^2$ .

In 2. and 3.,  $\text{ker}(T) = \{\mathbf{0}\}$  and  $\text{im}(T) = \mathbb{R}^2$ .

In 4.,  $\text{ker}(T)$  is the set of all constant polynomials (i.e. those of degree 0), and  $\text{im}(T) = \mathbb{R}[x]$ .

In 5.,  $\text{ker}(S_\alpha) = \{\mathbf{0}\}$ , and  $\text{im}(S_\alpha) = K[x]$ , while  $\text{ker}(E_\alpha)$  is the set of all polynomials divisible by  $x - \alpha$ , and  $\text{im}(E_\alpha) = K$ .

In 6.,  $\text{ker}(I_V) = \{\mathbf{0}\}$  and  $\text{im}(T) = V$ .

In 7.,  $\text{ker}(\mathbf{0}_{U,V}) = U$  and  $\text{im}(\mathbf{0}_{U,V}) = \{\mathbf{0}\}$ .

**Proposition 5.3** (i)  $\text{im}(T)$  is a subspace of  $V$ ; (ii)  $\text{ker}(T)$  is a subspace of  $U$ .

PROOF: (i) We must show that  $\text{im}(T)$  is closed under addition and scalar multiplication. Let  $\mathbf{v}_1, \mathbf{v}_2 \in \text{im}(T)$ . Then  $\mathbf{v}_1 = T(\mathbf{u}_1)$ ,  $\mathbf{v}_2 = T(\mathbf{u}_2)$  for some  $\mathbf{u}_1, \mathbf{u}_2 \in U$ . Then

$$\mathbf{v}_1 + \mathbf{v}_2 = T(\mathbf{u}_1) + T(\mathbf{u}_2) = T(\mathbf{u}_1 + \mathbf{u}_2) \in \text{im}(T); \quad \alpha \mathbf{v}_1 = \alpha T(\mathbf{u}_1) = T(\alpha \mathbf{u}_1) \in \text{im}(T),$$

so  $\text{im}(T)$  is a subspace of  $V$ .

(ii) Similarly, we must show that  $\ker(T)$  is closed under addition and scalar multiplication. Let  $\mathbf{u}_1, \mathbf{u}_2 \in \ker(T)$ . Then

$$T(\mathbf{u}_1 + \mathbf{u}_2) = T(\mathbf{0}_U + \mathbf{0}_U) = T(\mathbf{0}_U) = \mathbf{0}_V; \quad T(\alpha\mathbf{u}_1) = \alpha T(\mathbf{u}_1) = \alpha\mathbf{0}_V = \mathbf{0}_V,$$

so  $\mathbf{u}_1 + \mathbf{u}_2, \alpha\mathbf{u}_1 \in \ker(T)$  and  $\ker(T)$  is a subspace of  $U$ .  $\square$

### 5.3 Rank and nullity

#### Definition

- (i)  $\dim(\text{im}(T))$  is called the *rank* of  $T$ ;
- (ii)  $\dim(\ker(T))$  is called the *nullity* of  $T$ .

**Theorem 5.4** (The Dimension Theorem) *Let  $U, V$  be vector spaces over  $K$  with  $U$  finite-dimensional, and let  $T : U \rightarrow V$  be a linear map. Then*

$$\dim(\text{im}(T)) + \dim(\ker(T)) = \dim(U); \quad \text{i.e.} \quad \text{rank}(T) + \text{nullity}(T) = \dim(U).$$

PROOF: Since  $U$  is finite-dimensional and  $\ker(T)$  is a subspace of  $U$ ,  $\ker(T)$  is finite-dimensional. Let  $\text{nullity}(T) = s$  and let  $\mathbf{e}_1, \dots, \mathbf{e}_s$  be a basis of  $\ker(T)$ . By Theorem 3.6, we can extend  $\mathbf{e}_1, \dots, \mathbf{e}_s$  to a basis  $\mathbf{e}_1, \dots, \mathbf{e}_s, \mathbf{f}_1, \dots, \mathbf{f}_r$  of  $U$ . Then  $\dim(U) = s + r$ , so to prove the theorem we have to prove that  $\dim(\text{im}(T)) = r$ .

Clearly  $T(\mathbf{e}_1), \dots, T(\mathbf{e}_s), T(\mathbf{f}_1), \dots, T(\mathbf{f}_r)$  span  $\text{im}(T)$ , and since  $T(\mathbf{e}_1) = \dots = T(\mathbf{e}_s) = \mathbf{0}_V$  this implies that  $T(\mathbf{f}_1), \dots, T(\mathbf{f}_r)$  span  $\text{im}(T)$ . We shall show that  $T(\mathbf{f}_1), \dots, T(\mathbf{f}_r)$  are linearly independent.

Suppose that, for some scalars  $\alpha_i$ , we have

$$\alpha_1 T(\mathbf{f}_1) + \dots + \alpha_r T(\mathbf{f}_r) = \mathbf{0}_V.$$

Then  $T(\alpha_1 \mathbf{f}_1 + \dots + \alpha_r \mathbf{f}_r) = \mathbf{0}_V$ , so  $\alpha_1 \mathbf{f}_1 + \dots + \alpha_r \mathbf{f}_r \in \ker(T)$ . But  $\mathbf{e}_1, \dots, \mathbf{e}_s$  is a basis of  $\ker(T)$ , so there exist scalars  $\beta_i$  with

$$\alpha_1 \mathbf{f}_1 + \dots + \alpha_r \mathbf{f}_r = \beta_1 \mathbf{e}_1 + \dots + \beta_s \mathbf{e}_s \implies \alpha_1 \mathbf{f}_1 + \dots + \alpha_r \mathbf{f}_r - \beta_1 \mathbf{e}_1 - \dots - \beta_s \mathbf{e}_s = \mathbf{0}_U.$$

But we know that  $\mathbf{e}_1, \dots, \mathbf{e}_s, \mathbf{f}_1, \dots, \mathbf{f}_r$  form a basis of  $U$ , so they are linearly independent, and hence

$$\alpha_1 = \dots = \alpha_r = \beta_1 = \dots = \beta_s = 0,$$

and we have proved that  $T(\mathbf{f}_1), \dots, T(\mathbf{f}_r)$  are linearly independent.

Since  $T(\mathbf{f}_1), \dots, T(\mathbf{f}_r)$  both span  $\text{im}(T)$  and are linearly independent, they form a basis of  $\text{im}(U)$ , and hence  $\dim(\text{im}(T)) = r$ , which completes the proof.  $\square$

**Examples** Once again, we consider Examples 1 – 7 above. To deal with finite-dimensional spaces we restrict to an  $(n+1)$ -dimensional space  $K[x]_{\leq n}$  in Examples 4 and 5, that is, we consider  $T : \mathbb{R}[x]_{\leq n} \rightarrow \mathbb{R}[x]_{\leq n}$ ,  $S_\alpha : K[x]_{\leq n} \rightarrow K[x]_{\leq n}$ , and  $E_\alpha : K[x]_{\leq n} \rightarrow K$  correspondingly. Let  $n = \dim(U) = \dim(V)$  in 5. and 6.

Example	rank( $T$ )	nullity( $T$ )	dim( $U$ )
1.	2	1	3
2.	2	0	2
3.	2	0	2
4.	$n$	1	$n + 1$

Example	rank( $T$ )	nullity( $T$ )	dim( $U$ )
5. $S_\alpha$	$n + 1$	0	$n + 1$
5. $E_\alpha$	1	$n$	$n + 1$
6.	$n$	0	$n$
7.	0	$n$	$n$

**Corollary 5.5** Let  $T : U \rightarrow V$  be a linear map, where  $\dim(U) = \dim(V) = n$ . Then the following properties of  $T$  are equivalent:

- (i)  $T$  is surjective;
- (ii)  $\text{rank}(T) = n$ ;
- (iii)  $\text{nullity}(T) = 0$ ;
- (iv)  $T$  is injective;
- (v)  $T$  is bijective;

PROOF:  $T$  is surjective  $\Leftrightarrow \text{im}(T) = V$ , so clearly (i)  $\Rightarrow$  (ii). But if  $\text{rank}(T) = n$ , then  $\dim(\text{im}(T)) = \dim(V)$  so (by Corollary 3.9) a basis of  $\text{im}(T)$  is a basis of  $V$ , and hence  $\text{im}(T) = V$ . Thus (i)  $\Leftrightarrow$  (ii).

(ii)  $\Leftrightarrow$  (iii) follows directly from Theorem 5.4.

$\text{nullity}(T) = 0 \Leftrightarrow \ker(T) = \{\mathbf{0}\}$  so clearly (iv)  $\Rightarrow$  (iii). If  $\ker(T) = \{\mathbf{0}\}$  and  $T(\mathbf{u}_1) = T(\mathbf{u}_2)$  then  $T(\mathbf{u}_1 - \mathbf{u}_2) = \mathbf{0}$ , so  $\mathbf{u}_1 - \mathbf{u}_2 \in \ker(T) = \{\mathbf{0}\}$ , which implies  $\mathbf{u}_1 = \mathbf{u}_2$  and  $T$  is injective. Thus (iii)  $\Leftrightarrow$  (iv). (In fact, (iii)  $\Leftrightarrow$  (iv) is true for any linear map  $T$ .)

Finally, (v) is equivalent to (i) and (iv), which are equivalent to each other.  $\square$

**Definition** If the conditions in the above corollary are met, then  $T$  is called a *non-singular* linear map. Otherwise,  $T$  is called *singular*. Notice that the terms singular and non-singular are only used for linear maps  $T : U \rightarrow V$  for which  $U$  and  $V$  have the same dimension.

## 5.4 Operations on linear maps

We define the operations of *addition*, *scalar multiplication*, and *composition* on linear maps.

Let  $T_1 : U \rightarrow V$  and  $T_2 : U \rightarrow V$  be two linear maps, and let  $\alpha \in K$  be a scalar.

**Addition:** We define a map  $T_1 + T_2 : U \rightarrow V$  by the rule  $(T_1 + T_2)(\mathbf{u}) = T_1(\mathbf{u}) + T_2(\mathbf{u})$  for  $\mathbf{u} \in U$ .

**Scalar multiplication:** We define a map  $\alpha T_1 : U \rightarrow V$  by the rule  $(\alpha T_1)(\mathbf{u}) = \alpha T_1(\mathbf{u})$  for  $\mathbf{u} \in U$ .

Now let  $T_1 : U \rightarrow V$  and  $T_2 : V \rightarrow W$  be two linear maps.

**Composition:** We define a map  $T_2 T_1 : U \rightarrow W$  by  $(T_2 T_1)(\mathbf{u}) = T_2(T_1(\mathbf{u}))$  for  $\mathbf{u} \in U$ . In particular, we define  $T^2 = TT$  and  $T^{i+1} = T^i T$  for  $i > 2$ .

It is routine to check that  $T_1 + T_2$ ,  $\alpha T_1$  and  $T_2 T_1$  are themselves all linear maps.

Furthermore, for fixed vector spaces  $U$  and  $V$  over  $K$ , the operations of addition and scalar multiplication on the set  $\text{Hom}_K(U, V)$  of all linear maps from  $U$  to  $V$  makes  $\text{Hom}_K(U, V)$  into a vector space over  $K$ .

A vector space  $U^* = \text{Hom}_K(U, K)$  plays a special role. It is often called *the dual space* or *the space of covectors*. One can think of coordinates as elements of  $U^*$ . Indeed, let  $\mathbf{e}_i$  be a basis of  $U$ . Every  $\mathbf{x} \in U$  can be uniquely written as

$$\mathbf{x} = \alpha_1 \mathbf{e}_1 + \dots + \alpha_n \mathbf{e}_n, \quad \alpha_i \in K.$$

The elements  $\alpha_i$  depend on  $\mathbf{x}$  as well as on a choice of the basis, so for each  $i$  one can write the coordinate function

$$\mathbf{e}^i : U \rightarrow K, \quad \mathbf{e}^i(\mathbf{x}) = \alpha_i.$$

It is routine to check that  $\mathbf{e}^i$  is a linear map.

## 6 Matrices

The material in this section will be familiar to many of you already.

Let  $K$  be a field and  $m, n \in \mathbb{N}$ . An  $m \times n$  matrix  $A$  over  $K$  is an  $m \times n$  rectangular array of numbers (i.e. scalars) in  $K$ . The entry in row  $i$  and column  $j$  is usually written  $\alpha_{ij}$ . (We use the corresponding Greek letter.) We write  $A = (\alpha_{ij})$  to make things clear.

For example, we could take

$$K = \mathbb{R}, \quad m = 3, \quad n = 4, \quad A = \begin{pmatrix} 2 & -1 & -\pi & 0 \\ 3 & -3/2 & 0 & 6 \\ -1.23 & 0 & 10^{10} & 0 \end{pmatrix},$$

and then  $\alpha_{13} = -\pi$ ,  $\alpha_{33} = 10^{10}$ ,  $\alpha_{34} = 0$ , etc.

**Addition of matrices.** Let  $A = (\alpha_{ij})$  and  $B = (\beta_{ij})$  be two  $m \times n$  matrices over  $K$ . We define  $A + B$  to be the  $m \times n$  matrix  $C = (\gamma_{ij})$ , where  $\gamma_{ij} = \alpha_{ij} + \beta_{ij}$  for all  $i, j$ . For example,

$$\begin{pmatrix} 1 & 3 \\ 0 & 2 \end{pmatrix} + \begin{pmatrix} -2 & -3 \\ 1 & -4 \end{pmatrix} = \begin{pmatrix} -1 & -0 \\ 1 & -2 \end{pmatrix}.$$

**Scalar multiplication.** Let  $A = (\alpha_{ij})$  be an  $m \times n$  matrix over  $K$  and let  $\beta \in K$  be a scalar. We define the scalar multiple  $\beta A$  to be the  $m \times n$  matrix  $C = (\gamma_{ij})$ , where  $\gamma_{ij} = \beta\alpha_{ij}$  for all  $i, j$ .

**Multiplication of matrices.** Let  $A = (\alpha_{ij})$  be an  $l \times m$  matrix over  $K$  and let  $B = (\beta_{ij})$  be an  $m \times n$  matrix over  $K$ . The product  $AB$  is an  $l \times n$  matrix  $C = (\gamma_{ij})$  where, for  $1 \leq i \leq l$  and  $1 \leq j \leq n$ ,

$$\gamma_{ij} = \sum_{k=1}^m \alpha_{ik}\beta_{kj} = \alpha_{i1}\beta_{1j} + \alpha_{i2}\beta_{2j} + \cdots + \alpha_{im}\beta_{mj}.$$

It is essential that the number  $m$  of columns of  $A$  is equal to the number of rows of  $B$ . Otherwise  $AB$  makes no sense. If you are familiar with scalar products of vectors, note also that  $\gamma_{ij}$  is the scalar product of the  $i$ -th row of  $A$  with the  $j$ -th column of  $B$ .

For example, let

$$A = \begin{pmatrix} 2 & 3 & 4 \\ 1 & 6 & 2 \end{pmatrix}, \quad B = \begin{pmatrix} 2 & 6 \\ 3 & 2 \\ 1 & 9 \end{pmatrix}.$$

Then

$$AB = \begin{pmatrix} 2 \times 2 + 3 \times 3 + 4 \times 1 & 2 \times 6 + 3 \times 2 + 4 \times 9 \\ 1 \times 2 + 6 \times 3 + 2 \times 1 & 1 \times 6 + 6 \times 2 + 2 \times 9 \end{pmatrix} = \begin{pmatrix} 17 & 54 \\ 22 & 36 \end{pmatrix},$$

$$BA = \begin{pmatrix} 10 & 42 & 20 \\ 8 & 21 & 16 \\ 11 & 57 & 22 \end{pmatrix}.$$

Let  $C = \begin{pmatrix} 2 & 3 & 1 \\ 6 & 2 & 9 \end{pmatrix}$ . Then  $AC$  and  $CA$  are not defined.

Let  $D = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ . Then  $AD$  is not defined, but  $DA = \begin{pmatrix} 4 & 15 & 8 \\ 1 & 6 & 2 \end{pmatrix}$ .

**Proposition 6.1** *Matrices satisfy the following laws whenever the sums and products involved are defined:*

- (i)  $A + B = B + A$ ;
- (ii)  $(A + B)C = AC + BC$ ;
- (iii)  $C(A + B) = CA + CB$ ;
- (iv)  $(\lambda A)B = \lambda(AB) = A(\lambda B)$ ;
- (v)  $(AB)C = A(BC)$ .

PROOF: These are all routine checks that the entries of the left-hand-sides are equal to the corresponding entries on the right-hand-side. Let us do (v) as an example.

Let  $A$ ,  $B$  and  $C$  be  $l \times m$ ,  $m \times n$  and  $n \times p$  matrices, respectively. Then  $AB = D = (\delta_{ij})$  is an  $l \times n$  matrix with  $\delta_{ij} = \sum_{s=1}^m \alpha_{is}\beta_{sj}$ , and  $BC = E = (\varepsilon_{ij})$  is an  $m \times p$  matrix with  $\varepsilon_{ij} = \sum_{t=1}^n \beta_{it}\gamma_{tj}$ . Then  $(AB)C = DC$  and  $A(BC) = AE$  are both  $l \times p$  matrices, and we have to show that their coefficients are equal. The  $(i, j)$ -coefficient of  $DC$  is

$$\sum_{t=1}^n \delta_{it}\gamma_{tj} = \sum_{t=1}^n \left( \sum_{s=1}^m \alpha_{is}\beta_{st} \right) \gamma_{tj} = \sum_{s=1}^m \alpha_{is} \left( \sum_{t=1}^n \beta_{st}\gamma_{tj} \right) = \sum_{s=1}^m \alpha_{is}\varepsilon_{sj}$$

which is the  $(i, j)$ -coefficient of  $AE$ . Hence  $(AB)C = A(BC)$ .  $\square$

**The zero and identity matrices.** The  $m \times n$  zero matrix  $\mathbf{0}_{mn}$  over any field  $K$  has all of its entries equal to 0.

The  $n \times n$  identity matrix  $I_n = (\alpha_{ij})$  over any field  $K$  has  $\alpha_{ii} = 1$  for  $1 \leq i \leq n$ , but  $\alpha_{ij} = 0$  when  $i \neq j$ . For example,

$$I_1 = (1), \quad I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad I_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Note that  $I_n A = A$  for any  $n \times m$  matrix  $A$  and  $A I_n = A$  for any  $m \times n$  matrix  $A$ .

**Row and column vectors.** The set of all  $m \times n$  matrices over  $K$  will be denoted by  $K^{m,n}$ . Note that  $K^{m,n}$  is itself a vector space over  $K$  using the operations of addition and scalar multiplication defined above, and it has dimension  $mn$ . (This should be obvious - is it?)

A  $1 \times n$  matrix is called a *row vector*. We will regard  $K^{1,n}$  as being the same as  $K^n$ .

A  $n \times 1$  matrix is called a *column vector*. We will denote the the space  $K^{n,1}$  of all column vectors by  $K^{n,1}$ . In matrix calculations, we will use  $K^{n,1}$  more often than  $K^n$ .

## 7 Linear Transformations and Matrices

We shall see in this section that, for fixed choice of bases, there is a very natural one-one correspondence between linear maps and matrices, such that the operations on linear maps and matrices defined in Chapters 5 and 6 also correspond to each other. This is perhaps the most important idea in linear algebra, because it enables us to deduce properties of matrices from those of linear maps, and vice-versa.

### 7.1 Setting up the correspondence

Let  $T : U \rightarrow V$  be a linear map, where  $\dim(U) = n$ ,  $\dim(V) = m$ . Choose a basis  $\mathbf{e}_1, \dots, \mathbf{e}_n$  of  $U$  and a basis  $\mathbf{f}_1, \dots, \mathbf{f}_m$  of  $V$ .

Now, for  $1 \leq j \leq n$ ,  $T(\mathbf{e}_j) \in V$ , so  $T(\mathbf{e}_j)$  can be written uniquely as a linear combination of

$\mathbf{f}_1, \dots, \mathbf{f}_m$ . Let

$$\begin{aligned} T(\mathbf{e}_1) &= \alpha_{11}\mathbf{f}_1 + \alpha_{21}\mathbf{f}_2 + \cdots + \alpha_{m1}\mathbf{f}_m \\ T(\mathbf{e}_2) &= \alpha_{12}\mathbf{f}_1 + \alpha_{22}\mathbf{f}_2 + \cdots + \alpha_{m2}\mathbf{f}_m \\ &\dots \\ T(\mathbf{e}_n) &= \alpha_{1n}\mathbf{f}_1 + \alpha_{2n}\mathbf{f}_2 + \cdots + \alpha_{mn}\mathbf{f}_m \end{aligned}$$

where the coefficients  $\alpha_{ij} \in K$  (for  $1 \leq i \leq m$ ,  $1 \leq j \leq n$ ) are uniquely determined. In other words,

$$T(\mathbf{e}_j) = \sum_{i=1}^m \alpha_{ij}\mathbf{f}_i \quad \text{for } 1 \leq j \leq n.$$

The coefficients  $\alpha_{ij}$  form an  $m \times n$  matrix

$$A = \begin{pmatrix} \alpha_{11} & \alpha_{12} & \cdots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \cdots & \alpha_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ \alpha_{m1} & \alpha_{m2} & \cdots & \alpha_{mn} \end{pmatrix}$$

over  $K$ . Then  $A$  is called the matrix of the linear map  $T$  with respect to the chosen bases of  $U$  and  $V$ . In general, different choice of bases gives different matrices. We shall address this issue later in the course, in Section 11.)

Notice the role of individual columns in  $A$ . The  $j$ -th column of  $A$  consists of the coefficients of  $\mathbf{f}_i$  in  $T(\mathbf{e}_j)$ .

**Theorem 7.1** *Let  $U, V$  be vector spaces over  $K$  of dimensions  $n, m$ , respectively. Then, for a given choice of bases of  $U$  and  $V$ , there is a one-one correspondence between the set  $\text{Hom}_K(U, V)$  of linear maps  $U \rightarrow V$  and the set  $K^{m,n}$  of  $m \times n$  matrices over  $K$ .*

PROOF: As we saw above, any linear map  $T : U \rightarrow V$  determines an  $m \times n$  matrix  $A$  over  $K$ .

Conversely, let  $A = (\alpha_{ij})$  be an  $m \times n$  matrix over  $K$ . Then, by Proposition 5.2, there is just one linear  $T : U \rightarrow V$  with  $T(\mathbf{e}_j) = \sum_{i=1}^m \alpha_{ij}\mathbf{f}_i$  for  $1 \leq j \leq n$ , so we have a one-one correspondence.  $\square$

**Examples** Once again, we consider our examples from Section 5.

1.  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ ,  $T(\alpha, \beta, \gamma) = (\alpha, \beta)$ . Usually, we choose the standard bases of  $K^m$  and  $K^n$ , which in this case are  $\mathbf{e}_1 = (1, 0, 0)$ ,  $\mathbf{e}_2 = (0, 1, 0)$ ,  $\mathbf{e}_3 = (0, 0, 1)$  and  $\mathbf{f}_1 = (1, 0)$ ,  $\mathbf{f}_2 = (0, 1)$ . We have  $T(\mathbf{e}_1) = \mathbf{f}_1$ ,  $T(\mathbf{e}_2) = \mathbf{f}_2$ ,  $T(\mathbf{e}_3) = \mathbf{0}$ , and the matrix is

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

But suppose we chose a different basis, say  $\mathbf{e}_1 = (1, 1, 1)$ ,  $\mathbf{e}_2 = (0, 1, 1)$ ,  $\mathbf{e}_3 = (1, 0, 1)$ ,  $\mathbf{f}_1 = (0, 1)$ ,  $\mathbf{f}_2 = (1, 0)$ . Then we have  $T(\mathbf{e}_1) = (1, 1) = \mathbf{f}_1 + \mathbf{f}_2$ ,  $T(\mathbf{e}_2) = (0, 1) = \mathbf{f}_1$ ,  $T(\mathbf{e}_3) = (1, 0) = \mathbf{f}_2$ , and the matrix is

$$\begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}.$$

2.  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ ,  $T$  is a rotation through  $\theta$  anti-clockwise about the origin. We saw that  $T(1, 0) = (\cos \theta, \sin \theta)$  and  $T(0, 1) = (-\sin \theta, \cos \theta)$ , so the matrix using the standard bases is

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

3.  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ ,  $T$  is a reflection through the line through the origin making an angle  $\theta/2$  with the  $x$ -axis. We saw that  $T(1, 0) = (\cos \theta, \sin \theta)$  and  $T(0, 1) = (\sin \theta, -\cos \theta)$ , so the matrix using the standard bases is

$$\begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix}.$$

4. This time we take the differentiation map  $T$  from  $\mathbb{R}[x]_{\leq n}$  to  $\mathbb{R}[x]_{\leq n-1}$ . Then, with respect to the bases  $1, x, x^2, \dots, x^n$  and  $1, x, x^2, \dots, x^{n-1}$  of  $\mathbb{R}[x]_{\leq n}$  and  $\mathbb{R}[x]_{\leq n-1}$ , respectively, the matrix of  $T$  is

$$\begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 2 & \dots & 0 \\ 0 & 0 & 0 & 3 & \dots & 0 \\ 0 & 0 & 0 & \dots & n-1 & 0 \\ 0 & 0 & 0 & \dots & \dots & n \end{pmatrix}.$$

5. Let  $S_\alpha : K[x]_{\leq n} \rightarrow K[x]_{\leq n}$  be the shift. With respect to the basis  $1, x, x^2, \dots, x^n$  of  $K[x]_{\leq n}$ , we calculate  $S_\alpha(x^n) = (x - \alpha)^n$ . The binomial formula gives the matrix of  $S_\alpha$ ,

$$\begin{pmatrix} 1 & -\alpha & \alpha^2 & \dots & (-1)^n \alpha^n \\ 0 & 1 & -2\alpha & \dots & (-1)^{n-1} n \alpha^{n-1} \\ 0 & 0 & 1 & 3 & \dots & (-1)^{n-2} \frac{n(n-1)}{2} \alpha^{n-2} \\ \vdots & & & & \vdots & \\ 0 & 0 & 0 & \dots & 1 & -n\alpha \\ 0 & 0 & 0 & \dots & \dots & 1 \end{pmatrix}.$$

In the same basis of  $K[x]_{\leq n}$  and the basis  $1$  of  $K$ ,  $E_\alpha(x^n) = \alpha^n$ . The matrix of  $E_\alpha$  is

$$(1 \ \alpha \ \alpha^2 \ \dots \ \alpha^{n-1} \ \alpha^n)$$

6.  $T : V \rightarrow V$  is the identity map. Notice that  $U = V$  in this example. Provided that we choose the same basis for  $U$  and  $V$ , then the matrix of  $T$  is the  $n \times n$  identity matrix  $I_n$ . We shall be considering the situation where we use different bases for the domain and range of the identity map in Section 11.

7.  $T : U \rightarrow V$  is the zero map. The matrix of  $T$  is the  $m \times n$  zero matrix  $\mathbf{0}_{mn}$ .

For the given basis  $\mathbf{e}_1, \dots, \mathbf{e}_n$  of  $U$  and a vector  $\mathbf{u} = \lambda_1 \mathbf{e}_1 + \dots + \lambda_n \mathbf{e}_n \in U$ , let  $\underline{\mathbf{u}}$  denote the column vector

$$\begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \vdots \\ \lambda_n \end{pmatrix} \in K^{n,1},$$

and similarly, for the given basis  $\mathbf{f}_1, \dots, \mathbf{f}_m$  of  $V$  and a vector  $\mathbf{v} = \mu_1 \mathbf{f}_1 + \dots + \mu_m \mathbf{f}_m \in V$ , let  $\underline{\mathbf{v}}$  denote the column vector

$$\begin{pmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_m \end{pmatrix} \in K^{m,1}.$$

**Proposition 7.2** *Let  $T : U \rightarrow V$  be a linear map with matrix  $A = (\alpha_{ij})$ . Then  $T(\mathbf{u}) = \mathbf{v}$  if and only if  $\underline{\mathbf{A}}\underline{\mathbf{u}} = \underline{\mathbf{v}}$ .*

PROOF: We have

$$T(\mathbf{u}) = T\left(\sum_{j=1}^n \lambda_j \mathbf{e}_j\right) = \sum_{j=1}^n \lambda_j T(\mathbf{e}_j) = \sum_{j=1}^n \lambda_j \left(\sum_{i=1}^m \alpha_{ij} \mathbf{f}_i\right) = \sum_{i=1}^m \left(\sum_{j=1}^n \alpha_{ij} \lambda_j\right) \mathbf{f}_i = \sum_{i=1}^m \mu_i \mathbf{f}_i,$$

where  $\mu_i = \sum_{j=1}^n \alpha_{ij} \lambda_j$  is the entry in the  $i$ -th row of the column vector  $A\mathbf{u}$ . This proves the result.  $\square$

## 7.2 The correspondence between operations on linear maps and matrices

Let  $U$ ,  $V$  and  $W$  be vector spaces over the same field  $K$ , let  $\dim(U) = n$ ,  $\dim(V) = m$ ,  $\dim(W) = l$ , and choose fixed bases  $\mathbf{e}_1, \dots, \mathbf{e}_n$  of  $U$  and  $\mathbf{f}_1, \dots, \mathbf{f}_m$  of  $V$ , and  $\mathbf{g}_1, \dots, \mathbf{g}_l$  of  $W$ . All matrices of linear maps between these spaces will be written with respect to these bases.

**Addition.** Let  $T_1, T_2 : U \rightarrow V$  be linear maps with  $m \times n$  matrices  $A, B$  respectively. Then it is routine to check that the matrix of  $T_1 + T_2$  is  $A + B$ .

**Scalar multiplication.** Let  $T : U \rightarrow V$  be a linear map with  $m \times n$  matrices  $A$  and let  $\lambda \in K$  be a scalar. Then again it is routine to check that the matrix of  $\lambda T$  is  $\lambda A$ .

Note that the above two properties imply that the natural correspondence between linear maps and matrices is actually itself a linear map from  $\text{Hom}_K(U, V)$  to  $K^{m,n}$ .

**Composition of linear maps and matrix multiplication.** This time the correspondence is less obvious, and we state it as a theorem.

**Theorem 7.3** *Let  $T_1 : V \rightarrow W$  be a linear map with  $l \times m$  matrix  $A = (\alpha_{ij})$  and let  $T_2 : U \rightarrow V$  be a linear map with  $m \times n$  matrix  $B = (\beta_{ij})$ . Then the matrix of the composite map  $T_1 T_2 : U \rightarrow W$  is  $AB$ .*

PROOF: Let  $AB$  be the  $l \times n$  matrix  $(\gamma_{ij})$ . Then by the definition of matrix multiplication, we have  $\gamma_{ik} = \sum_{j=1}^m \alpha_{ij} \beta_{jk}$  for  $1 \leq i \leq l, 1 \leq k \leq n$ .

Let us calculate the matrix of  $T_1 T_2$ . We have

$$T_1 T_2(\mathbf{e}_k) = T_1\left(\sum_{j=1}^m \beta_{jk} \mathbf{f}_j\right) = \sum_{j=1}^m \beta_{jk} T_1(\mathbf{f}_j) = \sum_{j=1}^m \beta_{jk} \sum_{i=1}^l \alpha_{ij} \mathbf{g}_i = \sum_{i=1}^l \left(\sum_{j=1}^m \alpha_{ij} \beta_{jk}\right) \mathbf{g}_i = \sum_{i=1}^l \gamma_{ik} \mathbf{g}_i,$$

so the matrix of  $T_1 T_2$  is  $(\gamma_{ik}) = AB$  as claimed.  $\square$

### Examples

1. Let  $R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a rotation through an angle  $\theta$  anti-clockwise about the origin. We have seen that the matrix of  $R_\theta$  (using the standard basis) is  $\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ . Now clearly  $R_\theta$  followed by  $R_\phi$  is equal to  $R_{\theta+\phi}$ . We can check the corresponding result for matrices:

$$\begin{aligned} \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} &= \begin{pmatrix} \cos \phi \cos \theta - \sin \phi \sin \theta & -\cos \phi \sin \theta - \sin \phi \cos \theta \\ \sin \phi \cos \theta + \cos \phi \sin \theta & -\sin \phi \sin \theta + \cos \phi \cos \theta \end{pmatrix} \\ &= \begin{pmatrix} \cos(\phi + \theta) & -\sin(\phi + \theta) \\ \sin(\phi + \theta) & \cos(\phi + \theta) \end{pmatrix}. \end{aligned}$$

Note that in this case  $T_1 T_2 = T_2 T_1$ .

2. Let  $R_\theta$  be as in Example 1, and let  $M_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a reflection through a line through the origin at an angle  $\theta/2$  to the  $x$ -axis. We have seen that the matrix of  $M_\theta$  is

$\begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix}$ . What is the effect of doing first  $R_\theta$  and then  $M_\phi$ ? In this case, it might be easier (for some people) to work it out using the matrix multiplication! We have

$$\begin{aligned} \begin{pmatrix} \cos \phi & \sin \phi \\ \sin \phi & -\cos \phi \end{pmatrix} \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} &= \begin{pmatrix} \cos \phi \cos \theta + \sin \phi \sin \theta & -\cos \phi \sin \theta + \sin \phi \cos \theta \\ \sin \phi \cos \theta - \cos \phi \sin \theta & -\sin \phi \sin \theta - \cos \phi \cos \theta \end{pmatrix} \\ &= \begin{pmatrix} \cos(\phi - \theta) & \sin(\phi - \theta) \\ \sin(\phi - \theta) & -\cos(\phi - \theta) \end{pmatrix}, \end{aligned}$$

which is the matrix of  $M_{\phi-\theta}$ .

We get a different result if we do first  $M_\phi$  and then  $R_\theta$ . What do we get then?

### 7.3 Linear equations and the inverse image problem

The study and solution of systems of simultaneous linear equations is the main motivation behind the development of the theory of linear algebra and of matrix operations. Let us consider a system of  $m$  equations in  $n$  unknowns  $x_1, x_2 \dots x_n$ ,  $m, n \geq 1$ .

$$\begin{cases} \alpha_{11}x_1 + \alpha_{12}x_2 + \dots + \alpha_{1n}x_n = \beta_1 \\ \alpha_{21}x_1 + \alpha_{22}x_2 + \dots + \alpha_{2n}x_n = \beta_2 \\ \dots \\ \alpha_{m1}x_1 + \alpha_{m2}x_2 + \dots + \alpha_{mn}x_n = \beta_m \end{cases} \quad (1)$$

All coefficients  $\alpha_{ij}$  and  $\beta_i$  belong to  $K$ . Solving this system means finding all collections  $x_1, x_2 \dots x_n \in K$  such that the equations (1) hold.

Let  $A = (\alpha_{ij}) \in K^{m,n}$  be the  $m \times n$  matrix of coefficients. The crucial step is to introduce the column vectors

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \cdot \\ \cdot \\ x_n \end{pmatrix} \in K^{n,1} \quad \text{and} \quad \mathbf{b} = \begin{pmatrix} \beta_1 \\ \beta_2 \\ \cdot \\ \cdot \\ \beta_m \end{pmatrix} \in K^{m,1}.$$

This allows us to rewrite system (1) as a single equation

$$\mathbf{Ax} = \mathbf{b} \quad (2)$$

where the coefficient  $A$  is a matrix, the right hand side  $\mathbf{b}$  is a vector in  $K^{m,1}$  and the unknown  $\mathbf{x}$  is a vector  $K^{n,1}$ .

Using the notation of linear maps, we have just reduced solving a system of linear equations to the *inverse image* problem. That is, given a linear map  $T : U \rightarrow V$ , and a fixed vector  $\mathbf{v} \in V$ , find all  $\mathbf{u} \in U$  such that  $T(\mathbf{u}) = \mathbf{v}$ .

In fact, these two problems are equivalent! In the opposite direction, let us first forget all about  $A$ ,  $\mathbf{x}$  and  $\mathbf{b}$ . Then we choose bases in  $U$  and  $V$  and denote a matrix of  $T$  in these bases by  $A$ , the row vector of coordinates  $\mathbf{u}$  by  $\mathbf{x}$  and the row vector of coordinates  $\mathbf{v}$  by  $\mathbf{b}$ . Proposition 7.2 says that  $T(\mathbf{u}) = \mathbf{v}$  if and only if  $\mathbf{Ax} = \mathbf{b}$ . This reduces the inverse image problem to solving a system of linear equations.

Let us make several easy observations about the inverse image problem.

The case when  $\mathbf{v} = \mathbf{0}$  or, equivalently when  $\beta_i = 0$  for  $1 \leq i \leq m$ , is called the *homogeneous* case. Here the set of solutions is  $\{\mathbf{u} \in U \mid T(\mathbf{u}) = \mathbf{0}\}$ , which is precisely the kernel  $\ker(T)$  of  $T$ . The corresponding set of column vectors  $\mathbf{x} \in K^{n,1}$  with  $\mathbf{Ax} = \mathbf{0}$  is called the *nullspace* of the matrix  $A$ . So the nullity of  $A$  is the dimension of its nullspace.

In general, it is easy to see that if  $\mathbf{x}$  is one solution to a system of equations, then the complete set of solutions is equal to

$$\mathbf{x} + \text{nullspace}(A) = \{\mathbf{x} + \mathbf{y} \mid \mathbf{y} \in \text{nullspace}(A)\}.$$

It is possible that there are no solutions at all; this occurs when  $\mathbf{v} \notin \text{im}(T)$ . If there are solutions, then there is a unique solution precisely when  $\ker(T) = \{\mathbf{0}\}$ , or equivalently when  $\text{nullspace}(A) = \{\mathbf{0}\}$ . If the field  $K$  is infinite and there are solutions but  $\ker(T) \neq \{\mathbf{0}\}$ , then there are infinitely many solutions.

Now we would like to develop methods for solving the inverse image problem.

## 8 Elementary Operations and the Rank of a Matrix

### 8.1 Gauss transformations

There are two standard high school methods for solving linear systems: the *substitution method* (where you express variables in terms of the other variables and substitute the result in the remaining equations) and the *elimination method* (sometimes called the *Gauss method*). The latter is usually more effective, so we would like to contemplate its nature. Let us recall how it is done.

#### Examples

$$\begin{array}{rcl} 1. & 2x + y & = 1 & (1) \\ & 4x + 2y & = 1 & (2) \end{array}$$

Replacing (2) by (2)  $- 2 \times$  (1) gives  $0 = 1$ . This means that there are *no solutions*.

$$\begin{array}{rcl} 2. & 2x + y & = 1 & (1) \\ & 4x + y & = 1 & (2) \end{array}$$

Replacing (2) by (2)  $-$  (1) gives  $2x = 0$ . Replacing (1) by (1)  $- 2 \times$  (new 2) gives  $y = 1$ . Thus,  $(0, 1)$  is a *unique solution*.

$$\begin{array}{rcl} 3. & 2x + y & = 1 & (1) \\ & 4x + 2y & = 2 & (2) \end{array}$$

This time (2)  $-$  (1) gives  $0 = 0$ , so (2) is redundant.

After reduction, there is no equation with leading term  $y$ , which means that  $y$  can take on any value, say  $2\alpha$  (where the 2 is just to avoid fractions). The first equation determines  $x$  in terms of  $y$ , giving  $x = 1 - \alpha$ , so the general solution is  $(x, y) = (1 - \alpha, 2\alpha)$ ; there are *infinitely many solutions*.

Notice also that one solution is  $(x, y) = (1, 0)$ , and the general solution can be written as  $(x, y) = (1, 0) + \alpha(-1, 2)$ , where  $\alpha(-1, 2)$  is the solution of the corresponding *homogeneous system*  $2x + y = 0$ ;  $4x + 2y = 0$ .

$$\begin{array}{rcl} 4. & x + y + z & = 1 & (1) \\ & x + z & = 2 & (2) \\ & x - y + z & = 3 & (3) \\ & 3x + y + 3z & = 5 & (4) \end{array}$$

Now replacing (2) by (2)  $-$  (1) and then multiplying by  $-1$  gives  $y = -1$ . Replacing (3) by (3)  $-$  (1) gives  $-2y = 2$ , and replacing (4) by (4)  $- 3(1)$  also gives  $-2y = 2$ . So (3) and (4) both then reduce to  $0 = 0$ , and they are redundant.

$z$  does not occur as a leading term, so it can take any value, say  $\alpha$ , and then (2) gives  $y = -1$  and (1) gives  $x = 1 - y - z = 2 - \alpha$ , so the general solution is

$$(x, y, z) = (2 - \alpha, -1, \alpha) = (2, -1, 0) + \alpha(-1, 0, 1).$$

## 8.2 Elementary row operations

Many types of calculations with matrices can be carried out in a computationally efficient manner by the use of certain types of operations on rows and columns. We shall see later that these are really the same as the operations used in solving sets of simultaneous linear equations.

Let  $A$  be an  $m \times n$  matrix over  $K$  with rows  $\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_m \in K^{1,n}$ . The three types of elementary row operations on  $A$  are defined as follows.

(R1) For some  $i \neq j$ , add a multiple of  $\mathbf{r}_j$  to  $\mathbf{r}_i$ .

$$\text{Example: } \begin{pmatrix} 3 & 1 & 9 \\ 4 & 6 & 7 \\ 2 & 5 & 8 \end{pmatrix} \xrightarrow{\mathbf{r}_3 \rightarrow \mathbf{r}_3 - 3\mathbf{r}_1} \begin{pmatrix} 3 & 1 & 9 \\ 4 & 6 & 7 \\ -7 & 2 & -19 \end{pmatrix}$$

(R2) Interchange two rows

(R3) Multiply a row by a *non-zero* scalar.

$$\text{Example: } \begin{pmatrix} 2 & 0 & 5 \\ 1 & -2 & 3 \\ 5 & 1 & 2 \end{pmatrix} \xrightarrow{\mathbf{r}_2 \rightarrow 4\mathbf{r}_2} \begin{pmatrix} 2 & 0 & 5 \\ 4 & -8 & 12 \\ 5 & 1 & 2 \end{pmatrix}$$

## 8.3 Augmented Matrix

We would like to make the process of solving more mechanical by forgetting about the variable names  $w, x, y, z$ , etc. and doing the whole operation as a matrix calculation. For this, we use the *augmented matrix* of the system of equations, which for the system  $A\mathbf{x} = \underline{\beta}$  of  $m$  equations in  $n$  unknowns, where  $A$  is the  $m \times n$  matrix  $(\alpha_{ij})$  is defined to be the  $m \times (n+1)$  matrix

$$A = \left( \begin{array}{cccc|c} \alpha_{11} & \alpha_{12} & \dots & \alpha_{1n} & \beta_1 \\ \alpha_{21} & \alpha_{22} & \dots & \alpha_{2n} & \beta_2 \\ & & \dots & & \\ \alpha_{m1} & \alpha_{m2} & \dots & \alpha_{mn} & \beta_m \end{array} \right).$$

The vertical line in the matrix is put there just to remind us that the rightmost column is different from the others, and arises from the constants on the right hand side of the equations.

Let us look at the following system of linear equations over  $\mathbb{R}$ , that is, we want to find all  $w, x, y, z \in \mathbb{R}$  satisfying the equations.

$$\begin{cases} 2w - x + 4y - z = 1 \\ w + 2x + y + z = 2 \\ w - 3x + 3y - 2z = -1 \\ -3w - x - 5y = -3 \end{cases}$$

Elementary row operations on  $A$  are precisely Gauss transformations of the corresponding linear system. Thus, the solution can be carried out mechanically as follows:

Matrix	Operation
$\left( \begin{array}{cccc c} 2 & -1 & 4 & -1 & 1 \\ 1 & 2 & 1 & 1 & 2 \\ 1 & -3 & 3 & -2 & -1 \\ -3 & -1 & -5 & 0 & -3 \end{array} \right)$	$\mathbf{r}_1 \rightarrow \mathbf{r}_1/2$
$\left( \begin{array}{cccc c} 1 & -1/2 & 2 & -1/2 & 1/2 \\ 1 & 2 & 1 & 1 & 2 \\ 1 & -3 & 3 & -2 & -1 \\ -3 & -1 & -5 & 0 & -3 \end{array} \right)$	$\mathbf{r}_2 \rightarrow \mathbf{r}_2 - \mathbf{r}_1, \mathbf{r}_3 \rightarrow \mathbf{r}_3 - \mathbf{r}_1, \mathbf{r}_4 \rightarrow \mathbf{r}_4 + 3\mathbf{r}_1$
$\left( \begin{array}{cccc c} 1 & -1/2 & 2 & -1/2 & 1/2 \\ 0 & 5/2 & -1 & 3/2 & 3/2 \\ 0 & -5/2 & 1 & -3/2 & -3/2 \\ 0 & -5/2 & 1 & -3/2 & -3/2 \end{array} \right)$	$\mathbf{r}_3 \rightarrow \mathbf{r}_3 + \mathbf{r}_2, \mathbf{r}_4 \rightarrow \mathbf{r}_4 + \mathbf{r}_2$
$\left( \begin{array}{cccc c} 1 & -1/2 & 2 & -1/2 & 1/2 \\ 0 & 5/2 & -1 & 3/2 & 3/2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right)$	$\mathbf{r}_2 \rightarrow 2\mathbf{r}_2/5$
$\left( \begin{array}{cccc c} 1 & -1/2 & 2 & -1/2 & 1/2 \\ 0 & 1 & -2/5 & 3/5 & 3/5 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right)$	$\mathbf{r}_1 \rightarrow \mathbf{r}_1 + \mathbf{r}_2/2$
$\left( \begin{array}{cccc c} 1 & 0 & 9/5 & -1/5 & 4/5 \\ 0 & 1 & -2/5 & 3/5 & 3/5 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right)$	

The original system has been transformed to the following equivalent system, that is, both systems have the same solutions.

$$\begin{cases} w + 9y/5 - z/5 = 4/5 \\ x - 2y/5 + 3z/5 = 3/5 \end{cases}$$

In the latter system, variables  $y$  and  $z$  can take arbitrary values in  $\mathbb{R}$  in the solution set; say  $y = \alpha$ ,  $z = \beta$ . Then equations tell us that  $w = -9\alpha/5 + \beta/5 + 4/5$  and  $x = 2\alpha/5 - 3\beta/5 + 3/5$  (be careful to get the signs right!), and so the complete set of solutions is

$$\begin{aligned} (w, x, y, z) &= (-9\alpha/5 + \beta/5 + 4/5, 2\alpha/5 - 3\beta/5 + 3/5, \alpha, \beta) \\ &= (4/5, 3/5, 0, 0) + \alpha(-9/5, 2/5, 1, 0) + \beta(1/5, -3/5, 0, 1). \end{aligned}$$

#### 8.4 Row reduction a matrix

Let  $A = (\alpha_{ij})$  be an  $m \times n$  matrix over the field  $K$ . For the  $i$ -th row let  $\alpha_{i,c(i)}$  be the first (leftmost) non-zero entry in this row. In other words,  $\alpha_{i,c(i)} \neq 0$  while  $\alpha_{ij} = 0$  for all  $j < c(i)$ . By convention,  $c(i) = \infty$  if  $\alpha_{ij} = 0$  for all  $j$ .

After applying this procedure, the resulting matrix  $A = (\alpha_{ij})$  has the following properties.

- (i) All zero rows are below all non-zero rows.
- (ii) Let  $\mathbf{r}_1, \dots, \mathbf{r}_s$  be the non-zero rows. Then each  $\mathbf{r}_i$  with  $1 \leq i \leq s$  has 1 as its first non-zero entry. (In other words,  $\alpha_{i,c(i)} = 1$ .)
- (iii)  $c(1) < c(2) < \dots < c(s)$ .
- (iv)  $\alpha_{k,c(i)} = 0$  for all  $k > i$ .

**Definition** A matrix satisfying these properties is said to be in *upper echelon form*.

There is a stronger version of the last property

- (v)  $\alpha_{k,c(i)} = 0$  for all  $k \neq i$ .

**Definition** A matrix satisfying properties (i), (ii), (iii), and (v) is said to be in *row reduced form*.

An upper echelon form of a matrix will be used later to calculate the rank of a matrix. The row reduced form (the use of the definite article is intended: this form is, indeed, unique, though we shall not prove this) is used to solve systems of linear equations. In this light, the following theorem says that every system of linear equations can be solved by the Gauss (Elimination) method.

**Theorem 8.1** *Every matrix can be brought to row reduced form by elementary row transformations.*

PROOF: We describe an algorithm to achieve this. For a formal proof, we have to show:

- (i) after termination the resulting matrix has a row reduced form;
- (ii) the algorithm terminates after finitely many steps.

Both of these statements are clear from the nature of the algorithm. Make sure that you understand why they are clear!

At any stage in the procedure we are looking at the entry  $\alpha_{ij}$  in a particular position  $(i, j)$  of the matrix.  $(i, j)$  is called the *pivot* position, and  $\alpha_{ij}$  the *pivot* entry. We start with  $(i, j) = (1, 1)$  and proceed as follows.

1. If  $\alpha_{ij}$  and all entries below it in its columns are zero (i.e. if  $\alpha_{kj} = 0$  for all  $k \geq i$ ), then move the pivot one place to the right, to  $(i, j + 1)$  and repeat Step 1, or terminate if  $j = n$ .
2. If  $\alpha_{ij} = 0$  but  $\alpha_{kj} \neq 0$  for some  $k > i$  then apply (R2) and interchange  $\mathbf{r}_i$  and  $\mathbf{r}_k$ .
3. At this stage  $\alpha_{ij} \neq 0$ . If  $\alpha_{ij} \neq 1$ , then apply (R3) and multiply  $\mathbf{r}_i$  by  $\alpha_{ij}^{-1}$ .
4. At this stage  $\alpha_{ij} = 1$ . If, for any  $k \neq i$ ,  $\alpha_{kj} \neq 0$ , then apply (R1), and subtract  $\alpha_{kj}$  times  $\mathbf{r}_i$  from  $\mathbf{r}_k$ .
5. At this stage,  $\alpha_{kj} = 0$  for all  $k \neq i$ . If  $i = m$  or  $j = n$  then terminate. Otherwise, move the pivot diagonally down to the right to  $(i + 1, j + 1)$ , and go back to Step 1.

□

If one needs only an upper echelon form, this can done faster by replacing steps 4 and 5 with weaker and faster steps as follows.

- 4a. At this stage  $\alpha_{ij} = 1$ . If, for any  $k > i$ ,  $\alpha_{kj} \neq 0$ , then apply (R1), and subtract  $\alpha_{kj}$  times  $\mathbf{r}_i$  from  $\mathbf{r}_k$ .
- 5a. At this stage,  $\alpha_{kj} = 0$  for all  $k > i$ . If  $i = m$  or  $j = n$  then terminate. Otherwise, move the pivot diagonally down to the right to  $(i + 1, j + 1)$ , and go back to Step 1.

In the example below, we find an upper echelon form of a matrix by applying the faster

algorithm. The number in the ‘Step’ column refers to the number of the step applied in the description of the procedure above.

**Example**  $A = \begin{pmatrix} 0 & 0 & 1 & 2 & 1 \\ 2 & 4 & 2 & -4 & 2 \\ 3 & 6 & 3 & -6 & 3 \\ 1 & 2 & 3 & 3 & 3 \end{pmatrix}$ .

Matrix	Pivot	Step	Operation
$\begin{pmatrix} 0 & 0 & 1 & 2 & 1 \\ 2 & 4 & 2 & -4 & 2 \\ 3 & 6 & 3 & -6 & 3 \\ 1 & 2 & 3 & 3 & 3 \end{pmatrix}$	(1, 1)	<b>2</b>	$\mathbf{r}_1 \leftrightarrow \mathbf{r}_2$
$\begin{pmatrix} 2 & 4 & 2 & -4 & 2 \\ 0 & 0 & 1 & 2 & 1 \\ 3 & 6 & 3 & -6 & 3 \\ 1 & 2 & 3 & 3 & 3 \end{pmatrix}$	(1, 1)	<b>3</b>	$\mathbf{r}_1 \rightarrow \mathbf{r}_1/2$
$\begin{pmatrix} 1 & 2 & 1 & -2 & 1 \\ 0 & 0 & 1 & 2 & 1 \\ 3 & 6 & 3 & -6 & 3 \\ 1 & 2 & 3 & 3 & 3 \end{pmatrix}$	(1, 1)	<b>4</b>	$\mathbf{r}_3 \rightarrow \mathbf{r}_3 - 3\mathbf{r}_1$ $\mathbf{r}_4 \rightarrow \mathbf{r}_4 - \mathbf{r}_1$
$\begin{pmatrix} 1 & 2 & 1 & -2 & 1 \\ 0 & 0 & 1 & 2 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 5 & 2 \end{pmatrix}$	(1, 1) $\rightarrow$ (2, 2) $\rightarrow$ (2, 3)	<b>5, 1</b>	
$\begin{pmatrix} 1 & 2 & 1 & -2 & 1 \\ 0 & 0 & 1 & 2 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 5 & 2 \end{pmatrix}$	(2, 3)	<b>4</b>	$\mathbf{r}_4 \rightarrow \mathbf{r}_4 - 2\mathbf{r}_2$
Matrix	Pivot	Step	Operation
$\begin{pmatrix} 1 & 2 & 1 & -2 & 1 \\ 0 & 0 & 1 & 2 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}$	(2, 3) $\rightarrow$ (3, 4)	<b>5, 2</b>	$\mathbf{r}_3 \leftrightarrow \mathbf{r}_4$
$\begin{pmatrix} 1 & 2 & 1 & -2 & 1 \\ 0 & 0 & 1 & 2 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$	(3, 4) $\rightarrow$ (4, 5) $\rightarrow$ stop	<b>5, 1</b>	

### 8.5 Elementary column operations

In analogy to elementary row operations, one can introduce elementary column operations. Let  $A$  be an  $m \times n$  matrix over  $K$  with columns  $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$  as above. The three types of elementary column operations on  $A$  are defined as follows.

(C1) For some  $i \neq j$ , add a multiple of  $\mathbf{c}_j$  to  $\mathbf{c}_i$ .

Example:  $\begin{pmatrix} 3 & 1 & 9 \\ 4 & 6 & 7 \\ 2 & 5 & 8 \end{pmatrix} \xrightarrow{\mathbf{c}_3 \rightarrow \mathbf{c}_3 - 3\mathbf{c}_1} \begin{pmatrix} 3 & 1 & 0 \\ 4 & 6 & -5 \\ 2 & 5 & 2 \end{pmatrix}$

(C2) Interchange two columns.

(C3) Multiply a column by a *non-zero* scalar.

$$\text{Example: } \begin{pmatrix} 2 & 0 & 5 \\ 1 & -2 & 3 \\ 5 & 1 & 2 \end{pmatrix} \xrightarrow{\mathbf{c}_2 \rightarrow 4\mathbf{c}_2} \begin{pmatrix} 2 & 0 & 5 \\ 1 & -8 & 3 \\ 5 & 4 & 2 \end{pmatrix}$$

Elementary column operations change a linear system and cannot be applied to solve a system of linear equations. However, they are useful for reducing a matrix to a very nice form.

**Theorem 8.2** *By applying elementary row and column operations, a matrix can be brought into the block form*

$$\left( \begin{array}{c|c} I_s & \mathbf{0}_{s,n-s} \\ \hline \mathbf{0}_{m-s,s} & \mathbf{0}_{m-s,n-s} \end{array} \right),$$

where, as in Section 6,  $I_s$  denotes the  $s \times s$  identity matrix, and  $\mathbf{0}_{kl}$  the  $k \times l$  zero matrix.

PROOF: First, use elementary row operations to reduce  $A$  to row reduced form.

Now all  $\alpha_{i,c(i)} = 1$ . For each  $\alpha_{ij} \neq 0$  with  $j \neq c(i)$ , replace  $\mathbf{c}_j$  with  $\mathbf{c}_j - \alpha_{ij}\mathbf{c}_{c(i)}$ .

Finally the only nonzero entries of our matrix are  $\alpha_{i,c(i)} = 1$ . Now for each number  $i$  starting from  $i = 1$ , exchange  $\mathbf{c}_i$  and  $\mathbf{c}_{c(i)}$ .  $\square$

**Definition** The matrix in Theorem 8.2 is said to be in *row and column reduced form*, which is more usually called *Smith normal form*.

Let us look at an example of the second stage of procedure, that is, after reducing the matrix to the row reduced form.

Matrix	Operation
$\begin{pmatrix} 1 & 2 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$	$\mathbf{c}_2 \rightarrow \mathbf{c}_2 - 2\mathbf{c}_1$ $\mathbf{c}_5 \rightarrow \mathbf{c}_5 - \mathbf{c}_1$
$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$	$\mathbf{c}_2 \leftrightarrow \mathbf{c}_3$ $\mathbf{c}_5 \rightarrow \mathbf{c}_5 - 3\mathbf{c}_4$
$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 2 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$	$\mathbf{c}_3 \leftrightarrow \mathbf{c}_4$ $\mathbf{c}_5 \rightarrow \mathbf{c}_5 - 2\mathbf{c}_2$
$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$	

Now we would like to discuss the number  $s$  that appears in Theorem 8.2. Does the initial matrix uniquely determine this number?

## 8.6 The rank of a matrix

Let  $T : U \rightarrow V$  be a linear map, where  $\dim(U) = n$ ,  $\dim(V) = m$ . Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  be a basis of  $U$  and let  $\mathbf{f}_1, \dots, \mathbf{f}_m$  be a basis of  $V$ .

Recall from Section 5.3 that  $\text{rank}(T) = \dim(\text{im}(T))$ .

Now  $\text{im}(T)$  is spanned by the vectors  $T(\mathbf{e}_1), \dots, T(\mathbf{e}_n)$ , and by Theorem 3.5, some subset of these vectors forms a basis of  $\text{im}(T)$ . By definition of basis, this subset has size  $\dim(\text{im}(T)) = \text{rank}(T)$ , and by Corollary 3.9 no larger subset of  $T(\mathbf{e}_1), \dots, T(\mathbf{e}_n)$  can be linearly independent. We have therefore proved:

**Lemma 8.3**  $\text{rank}(T)$  is the size of the largest linearly independent subset of  $T(\mathbf{e}_1), \dots, T(\mathbf{e}_n)$ .

Now let  $A$  be an  $m \times n$  matrix over  $K$ . We shall denote the  $m$  rows of  $A$ , which are row vectors in  $K^n$  by  $\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_m$ , and similarly, we denote the  $n$  columns of  $A$ , which are column vectors in  $K^{m,1}$  by  $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$ .

**Definition**

(i) The *row-space* of  $A$  is the subspace of  $K^n$  spanned by the rows  $\mathbf{r}_1, \dots, \mathbf{r}_m$  of  $A$ . The *row rank* of  $A$  is equal to the dimension of the row-space of  $A$ . Equivalently, by the argument above, the row rank of  $A$  is equal to the size of the largest linearly independent subset of  $\mathbf{r}_1, \dots, \mathbf{r}_m$ .

(ii) The *column-space* of  $A$  is the subspace of  $K^{m,1}$  spanned by the columns  $\mathbf{c}_1, \dots, \mathbf{c}_n$  of  $A$ . The *column rank* of  $A$  is equal to the dimension of the column-space of  $A$ . Equivalently, by the argument above, the column rank of  $A$  is equal to the size of the largest linearly independent subset of  $\mathbf{c}_1, \dots, \mathbf{c}_n$ .

There is no obvious reason why there should be any particular relationship between the row and column ranks, but in fact it will turn out that they are always equal. First we show that the column rank is the same as the rank of the associated linear map.

**Theorem 8.4** Suppose that the linear map  $T$  has matrix  $A$ . Then  $\text{rank}(T)$  is equal to the column rank of  $A$ .

PROOF: As we saw in Section 7.1, the columns  $\mathbf{c}_1, \dots, \mathbf{c}_n$  of  $A$  are precisely the vectors  $T(\mathbf{e}_1), \dots, T(\mathbf{e}_n)$  written as column vectors. The result now follows directly from Lemma 8.3.  $\square$

**Example**

$$A = \begin{pmatrix} 1 & 2 & 0 & 1 & 1 \\ 2 & 4 & 1 & 3 & 0 \\ 4 & 8 & 0 & 4 & 4 \end{pmatrix} \begin{matrix} \mathbf{r}_1 \\ \mathbf{r}_2 \\ \mathbf{r}_3 \\ \mathbf{c}_1 & \mathbf{c}_2 & \mathbf{c}_3 & \mathbf{c}_4 & \mathbf{c}_5 \end{matrix}$$

We can calculate the row and column ranks by applying the sifting process (described in Section 3) to the row and column vectors, respectively.

Doing rows first,  $\mathbf{r}_1$  and  $\mathbf{r}_2$  are linearly independent, but  $\mathbf{r}_3 = 4\mathbf{r}_1$ , so the row rank is 2.

Now doing columns,  $\mathbf{c}_2 = 2\mathbf{c}_1$ ,  $\mathbf{c}_4 = \mathbf{c}_1 + \mathbf{c}_3$  and  $\mathbf{c}_5 = \mathbf{c}_1 - 2\mathbf{c}_3$ , so the column rank is also 2.

**Theorem 8.5** Applying (R1), (R2) or (R3) to a matrix  $A$  does not change the row or column rank of  $A$ . The same is true for (C1), (C2) and (C3).

PROOF: The row rank of  $A$  is the dimension of the row-space of  $A$ , which is the space of linear combinations  $\lambda_1\mathbf{r}_1 + \dots + \lambda_m\mathbf{r}_m$  of the rows of  $A$ . It is easy to see that (R1), (R2) and (R3) do not change this space, so they do not change the row-rank. (But notice that the scalar in (R3) must be non-zero for this to be true!)

The column rank of  $A = (\alpha_{ij})$  is the size of the largest linearly independent subset of  $\mathbf{c}_1, \dots, \mathbf{c}_n$ . Let  $\{\mathbf{c}_1, \dots, \mathbf{c}_s\}$  be some subset of the set  $\{\mathbf{c}_1, \dots, \mathbf{c}_n\}$  of columns of  $A$ . (We have written this as though the subset consisted of the first  $s$  columns, but this is just to keep the notation simple; it could be any subset of the columns.)

Then  $\mathbf{c}_1, \dots, \mathbf{c}_s$  are linearly dependent if and only if there exist scalars  $x_1, \dots, x_s \in K$ , not all zero, such that  $x_1\mathbf{c}_1 + x_2\mathbf{c}_2 + \dots + x_s\mathbf{c}_s = \mathbf{0}$ . If we write out the  $m$  components of this vector equation, we get a system of  $m$  simultaneous linear equations in the scalars  $x_i$  (which is why we have suddenly decided to call the scalars  $x_i$  rather than  $\lambda_i$ ).

$$\begin{aligned}\alpha_{11}x_1 + \alpha_{12}x_2 + \dots + \alpha_{1s}x_s &= 0 \\ \alpha_{21}x_1 + \alpha_{22}x_2 + \dots + \alpha_{2s}x_s &= 0 \\ &\dots \\ \alpha_{m1}x_1 + \alpha_{m2}x_2 + \dots + \alpha_{ms}x_s &= 0\end{aligned}$$

Now if we perform (R1), (R2) or (R3) on  $A$ , then we perform the corresponding operation on this system of equations. That is, we add a multiple of one equation to another, we interchange two equations, or we multiply one equation by a non-zero scalar. None of these operations change the set of solutions of the equations. Hence if they have some solution with the  $x_i$  not all zero before the operation, then they have the same solution after the operation. In other words, the elementary row operations do not change the linear dependence or independence of the set of columns  $\{\mathbf{c}_1, \dots, \mathbf{c}_s\}$ . Thus they do not change the size of the largest linearly independent subset of  $\mathbf{c}_1, \dots, \mathbf{c}_n$ , so they do not change the column rank of  $A$ .

The proof for the column operations (C1), (C2) and (C3) is the same with rows and columns interchanged.  $\square$

**Corollary 8.6** *Let  $s$  be the number of non-zero rows in the Smith normal form of a matrix  $A$  (see Theorem 8.2). Then both row rank of  $A$  and column rank of  $A$  are equal to  $s$ .*

PROOF: Since elementary operations preserve ranks, it suffices to find both ranks of a matrix in Smith normal form. But it is easy to see that the row space is precisely the space spanned by the first  $s$  standard vectors and hence has dimension  $s$ . Similarly the column space has dimension  $s$ .  $\square$

In particular, Corollary 8.6 establishes that the row rank is always equal to the column rank. This allows us to forget this distinction. From now we shall just talk about *the rank of a matrix*.

**Corollary 8.7** *The rank of a matrix  $A$  is equal to the number of non-zero rows after reducing  $A$  to upper echelon form.*

PROOF: The corollary follows from the fact that non-zero rows of a matrix in upper echelon form are linearly independent.

To see this, let  $\mathbf{r}_1, \dots, \mathbf{r}_s$  be the non-zero rows, and suppose that  $\lambda_1\mathbf{r}_1 + \dots + \lambda_s\mathbf{r}_s = \mathbf{0}$ . Now  $\mathbf{r}_1$  is the only row with a non-zero entry in column  $c(1)$ , so the entry in column  $c(1)$  of the vector  $\lambda_1\mathbf{r}_1 + \dots + \lambda_s\mathbf{r}_s$  is  $\lambda_1$ , and hence  $\lambda_1 = 0$ .

But then  $\mathbf{r}_2$  is the only row  $\mathbf{r}_k$  with  $k \geq 2$  with a non-zero entry in column  $c(2)$  and so the entry in column  $c(2)$  of the vector  $\lambda_2\mathbf{r}_2 + \dots + \lambda_s\mathbf{r}_s$  is  $\lambda_2$ , and hence  $\lambda_2 = 0$ . Continuing in this way (by induction), we find that  $\lambda_1 = \lambda_2 = \dots = \lambda_s = 0$ , and so  $\mathbf{r}_1, \dots, \mathbf{r}_s$  are linearly independent, as claimed.  $\square$

Corollary 8.7 give the most efficient way of computing the rank of a matrix. For instance, let us look at  $A = \begin{pmatrix} 1 & 2 & 0 & 1 & 1 \\ 2 & 4 & 1 & 3 & 0 \\ 4 & 8 & 1 & 5 & 2 \end{pmatrix}$ .

Matrix	Operation
$\begin{pmatrix} 1 & 2 & 0 & 1 & 1 \\ 2 & 4 & 1 & 3 & 0 \\ 4 & 8 & 1 & 5 & 2 \end{pmatrix}$	$\mathbf{r}_2 \rightarrow \mathbf{r}_2 - 2\mathbf{r}_1$ $\mathbf{r}_3 \rightarrow \mathbf{r}_3 - 4\mathbf{r}_1$
$\begin{pmatrix} 1 & 2 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & -2 \\ 0 & 0 & 1 & 1 & -2 \end{pmatrix}$	$\mathbf{r}_3 \rightarrow \mathbf{r}_3 - \mathbf{r}_2$
$\begin{pmatrix} 1 & 2 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$	

Since the resulting matrix in upper echelon form has 2 nonzero rows,  $\text{rank}(A) = 2$ .

## 8.7 Rank Criterion

The following theorem is proved in Assignment Sheet 6.

**Theorem 8.8** *Let  $A$  be the augmented  $n \times (m + 1)$  matrix of a linear system. Let  $B$  be the  $n \times m$  matrix obtained from  $A$  by removing the last column. The system of linear equations has a solution if and only if  $\text{rank}(A) = \text{rank}(B)$ .*

# 9 The Inverse of a Linear Transformation and of a Matrix

## 9.1 Definitions

As usual, let  $T : U \rightarrow V$  be a linear map with corresponding  $m \times n$  matrix  $A$ . If there is a map  $T^{-1} : V \rightarrow U$  with  $TT^{-1} = I_V$  and  $T^{-1}T = I_U$  then  $T$  is said to be *invertible*, and  $T^{-1}$  is called the *inverse* of  $T$ .

If this is the case, and  $A^{-1}$  is the  $(n \times m)$  matrix of  $T^{-1}$ , then we have  $AA^{-1} = I_m$  and  $A^{-1}A = I_n$ . We call  $A^{-1}$  the inverse of the matrix  $A$ , and say that  $A$  is invertible. Conversely, if  $A^{-1}$  is an  $n \times m$  matrix satisfying  $AA^{-1} = I_m$  and  $A^{-1}A = I_n$ , then the corresponding linear map  $T^{-1}$  satisfies  $TT^{-1} = I_V$  and  $T^{-1}T = I_U$ , so it is the inverse of  $T$ .

**Lemma 9.1** *Let  $A$  be a matrix of a linear map  $T$ . A linear map  $T$  is invertible if and only if its matrix  $A$  is invertible. The inverses  $T^{-1}$  and  $A^{-1}$  are unique.*

PROOF: It is clear that under the bijection between matrices and linear maps, invertible matrices correspond to invertible linear maps. This establishes the first statement.

Since the inverse map of a bijection is unique,  $T^{-1}$  is unique. Under the bijection between matrices and linear maps,  $A^{-1}$  must be the matrix of  $T^{-1}$ . Thus,  $A^{-1}$  is unique as well.  $\square$

**Theorem 9.2** *A linear map  $T$  is invertible if and only if  $T$  is non-singular. In particular, if  $T$  is invertible then  $m = n$ , so only square matrices can be invertible.*

**Remark** See Corollary 5.5 for the definition of non-singular linear maps. We may also say that the matrix  $A$  is non-singular if  $T$  is; but by this theorem, this is equivalent to  $A$  being invertible.

PROOF: If any map  $T$  has a left and right inverse, then it must be a bijection. Hence  $\ker(T) = \{\mathbf{0}\}$  and  $\text{im}(T) = V$ , so  $\text{nullity}(T) = 0$  and  $\text{rank}(T) = \dim(V) = m$ . But by Theorem 5.4, we have

$$n = \dim(U) = \text{rank}(T) + \text{nullity}(T) = m + 0 = m$$

and we see from the definition that  $T$  is non-singular.

Conversely, if  $n = m$  and  $T$  is non-singular, then by Corollary 5.5  $T$  is a bijection, and so it has an inverse  $T^{-1} : V \rightarrow U$  as a map. However, we still have to show that  $T^{-1}$  is a linear map. Let  $\mathbf{v}_1, \mathbf{v}_2 \in V$ . Then there exist  $\mathbf{u}_1, \mathbf{u}_2 \in U$  with  $T(\mathbf{u}_1) = \mathbf{v}_1$ ,  $T(\mathbf{u}_2) = \mathbf{v}_2$ . So  $T(\mathbf{u}_1 + \mathbf{u}_2) = \mathbf{v}_1 + \mathbf{v}_2$  and hence  $T^{-1}(\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u}_1 + \mathbf{u}_2$ . If  $\alpha \in K$ , then

$$T^{-1}(\alpha\mathbf{v}_1) = T^{-1}(T(\alpha\mathbf{u}_1)) = \alpha\mathbf{u}_1 = \alpha T^{-1}(\mathbf{v}_1),$$

so  $T^{-1}$  is linear, which completes the proof.  $\square$

**Example** Let  $A = \begin{pmatrix} 1 & 2 & 0 \\ 2 & 0 & 1 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & -2 \\ 0 & 1 \\ -2 & 5 \end{pmatrix}$ . Then  $AB = I_2$ , but  $BA \neq I_3$ ,

so a non-square matrix can have a right inverse which is not a left inverse. However, it can be deduced from Corollary 5.5 that if  $A$  is a square  $n \times n$  matrix and  $AB = I_n$  then  $A$  is non-singular, and then by multiplying  $AB = I_n$  on the left by  $A^{-1}$ , we see that  $B = A^{-1}$  and so  $BA = I_n$ .

This technique of multiplying on the left or right by  $A^{-1}$  is often used for transforming matrix equations. If  $A$  is invertible, then  $AX = B \iff X = A^{-1}B$  and  $XA = B \iff X = BA^{-1}$ .

**Lemma 9.3** *If  $A$  and  $B$  are invertible  $n \times n$  matrices, then  $AB$  is invertible, and  $(AB)^{-1} = B^{-1}A^{-1}$ .*

PROOF: This is clear, because  $ABB^{-1}A^{-1} = B^{-1}A^{-1}AB = I_n$ .  $\square$

## 9.2 Matrix inversion by row reduction

Two methods for finding the inverse of a matrix will be studied in this course. The first, using row reduction, which we shall look at now, is an efficient practical method similar to that used by computer packages. The second, using determinants, is of more theoretical interest, and will be done later in Section 10.

First note that if an  $n \times n$  matrix  $A$  is invertible, then it has rank  $n$ . Consider the row reduced form  $B = (\beta_{ij})$  of  $A$ . As we saw in Section 8.6, we have  $\beta_{ic(i)} = 1$  for  $1 \leq i \leq n$  (since  $\text{rank}(A) = \text{rank}(B) = n$ ), where  $c(1) < c(2) < \dots < c(n)$ , and clearly this is only possible if  $c(i) = i$  for  $1 \leq i \leq n$ . Then, since all other entries in column  $c(i)$  are zero, we have  $B = I_n$ . We have therefore proved:

**Proposition 9.4** *The row reduced form of an invertible  $n \times n$  matrix  $A$  is  $I_n$ .*

To compute  $A^{-1}$ , we reduce  $A$  to its row reduced form  $I_n$ , using elementary row operations, while simultaneously applying the same row operations, but starting with the identity matrix  $I_n$ . It turns out that these operations transform  $I_n$  to  $A^{-1}$ .

In practice, we might not know whether or not  $A$  is invertible before we start, but we will find out while carrying out this procedure because, if  $A$  is not invertible, then its rank will be less than  $n$ , and it will not row reduce to  $I_n$ .

First we will do an example to demonstrate the method, and then we will explain why it works. In the table below, the row operations applied are given in the middle column. The results of applying them to the matrix

$$A = \begin{pmatrix} 3 & 2 & 1 \\ 4 & 1 & 3 \\ 2 & 1 & 6 \end{pmatrix}$$

are given in the left column, and the results of applying them to  $I_3$  in the right column. So  $A^{-1}$  should be the final matrix in the right column.

$$\begin{array}{ccc} \begin{pmatrix} 3 & 2 & 1 \\ 4 & 1 & 3 \\ 2 & 1 & 6 \end{pmatrix} & & \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ & \mathbf{r}_1 \rightarrow \mathbf{r}_1/3 & \\ \begin{pmatrix} 1 & 2/3 & 1/3 \\ 4 & 1 & 3 \\ 2 & 1 & 6 \end{pmatrix} & & \begin{pmatrix} 1/3 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ & \mathbf{r}_2 \rightarrow \mathbf{r}_2 - 4\mathbf{r}_1 \\ & \mathbf{r}_3 \rightarrow \mathbf{r}_3 - 2\mathbf{r}_1 & \\ \begin{pmatrix} 1 & 2/3 & 1/3 \\ 0 & -5/3 & 5/3 \\ 0 & -1/3 & 16/3 \end{pmatrix} & & \begin{pmatrix} 1/3 & 0 & 0 \\ -4/3 & 1 & 0 \\ -2/3 & 0 & 1 \end{pmatrix} \\ & \mathbf{r}_2 \rightarrow -3\mathbf{r}_2/5 & \\ \begin{pmatrix} 1 & 2/3 & 1/3 \\ 0 & 1 & -1 \\ 0 & -1/3 & 16/3 \end{pmatrix} & & \begin{pmatrix} 1/3 & 0 & 0 \\ 4/5 & -3/5 & 0 \\ -2/3 & 0 & 1 \end{pmatrix} \\ & \mathbf{r}_1 \rightarrow \mathbf{r}_1 - 2\mathbf{r}_2/3 \\ & \mathbf{r}_3 \rightarrow \mathbf{r}_3 + \mathbf{r}_2/3 & \\ \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 5 \end{pmatrix} & & \begin{pmatrix} -1/5 & 2/5 & 0 \\ 4/5 & -3/5 & 0 \\ -2/5 & -1/5 & 1 \end{pmatrix} \\ & \mathbf{r}_3 \rightarrow \mathbf{r}_3/5 & \\ \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix} & & \begin{pmatrix} -1/5 & 2/5 & 0 \\ 4/5 & -3/5 & 0 \\ -2/25 & -1/25 & 1/5 \end{pmatrix} \\ & \mathbf{r}_1 \rightarrow \mathbf{r}_1 - \mathbf{r}_3 \\ & \mathbf{r}_2 \rightarrow \mathbf{r}_2 + \mathbf{r}_3 & \\ \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} & & \begin{pmatrix} -3/25 & 11/25 & -1/5 \\ 18/25 & -16/25 & 1/5 \\ -2/25 & -1/25 & 1/5 \end{pmatrix} \end{array}$$

So

$$A^{-1} = \begin{pmatrix} -3/25 & 11/25 & -1/5 \\ 18/25 & -16/25 & 1/5 \\ -2/25 & -1/25 & 1/5 \end{pmatrix}.$$

It is always a good idea to check the result afterwards. This is easier if we remove the common denominator 25, and we can then easily check that

$$\begin{pmatrix} 3 & 2 & 1 \\ 4 & 1 & 3 \\ 2 & 1 & 6 \end{pmatrix} \begin{pmatrix} -3 & 11 & -5 \\ 18 & -16 & 5 \\ -2 & -1 & 5 \end{pmatrix} = \begin{pmatrix} -3 & 11 & -5 \\ 18 & -16 & 5 \\ -2 & -1 & 5 \end{pmatrix} \begin{pmatrix} 3 & 2 & 1 \\ 4 & 1 & 3 \\ 2 & 1 & 6 \end{pmatrix} = \begin{pmatrix} 25 & 0 & 0 \\ 0 & 25 & 0 \\ 0 & 0 & 25 \end{pmatrix}$$

which confirms the result!

### 9.3 Elementary matrices

We shall now explain why the above method of calculating the inverse of a matrix works. Each elementary row operation on a matrix can be achieved by multiplying the matrix on the left by a corresponding matrix known as an *elementary matrix*. There are three types of these, all being slightly different from the identity matrix.

1.  $E(n)_{\lambda,i,j}^1$  (where  $i \neq j$ ) is the an  $n \times n$  matrix equal to the identity, but with an additional non-zero entry  $\lambda$  in the  $(i, j)$  position.
2.  $E(n)_{i,j}^2$  is the  $n \times n$  identity matrix with its  $i$ -th and  $j$ -th rows interchanged.
3.  $E(n)_{\lambda,i}^3$  (where  $\lambda \neq 0$ ) is the  $n \times n$  identity matrix with its  $(i, i)$  entry replaced by  $\lambda$ .

#### Examples

$$E(3)_{\frac{1}{3},1,3}^1 = \begin{pmatrix} 1 & 0 & \frac{1}{3} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad E(4)_{2,4}^2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad E(3)_{-4,3}^3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -4 \end{pmatrix}.$$

Let  $A$  be any  $m \times n$  matrix. Then  $E(m)_{\lambda,i,j}^1 A$  is the result we get by adding  $\lambda$  times the  $j$ -th row of  $A$  to the  $i$ -th row of  $A$ . Similarly  $E(m)_{i,j}^2 A$  is equal to  $A$  with its  $i$ -th and  $j$ -th rows interchanged, and  $E(m)_{\lambda,i}^3$  is equal to  $A$  with its  $i$ -th row multiplied by  $\lambda$ . You need to work out a few examples to convince yourself that this is true. For example

$$E(4)_{-2,4,2}^1 \begin{pmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 2 & 2 \\ 3 & 3 & 3 & 3 \\ 4 & 4 & 4 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -2 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 2 & 2 \\ 3 & 3 & 3 & 3 \\ 4 & 4 & 4 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 2 & 2 \\ 3 & 3 & 3 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

So, in the matrix inversion procedure, the effect of applying elementary row operations to reduce  $A$  to the identity matrix  $I_n$  is equivalent to multiplying  $A$  on the left by a sequence of elementary matrices. In other words, we have  $E_r E_{r-1} \dots E_1 A = I_n$ , for certain elementary  $n \times n$  matrices  $E_1, \dots, E_r$ . Hence  $E_r E_{r-1} \dots E_1 = A^{-1}$ . But when we apply the same elementary row operations to  $I_n$ , then we end up with  $E_r E_{r-1} \dots E_1 I_n = A^{-1}$ . This explains why the method works.

Notice also that the inverse of an elementary row matrix is another one of the same type. In fact it is easily checked that the inverses of  $E(n)_{\lambda,i,j}^1$ ,  $E(n)_{i,j}^2$  and  $E(n)_{\lambda,i}^3$  are respectively  $E(n)_{-\lambda,i,j}^1$ ,  $E(n)_{i,j}^2$  and  $E(n)_{\lambda^{-1},i}^3$ . Hence, if  $E_r E_{r-1} \dots E_1 A = I_n$  as in the preceding paragraph, then by using Lemma 9.3 we find that

$$A = (E_r E_{r-1} \dots E_1)^{-1} = E_1^{-1} E_2^{-1} \dots E_r^{-1},$$

which is itself a product of elementary matrices. We have proved:

**Theorem 9.5** *An invertible matrix is a product of elementary matrices.*

## 9.4 Application to linear equations

The most familiar examples of simultaneous equations are those where we have the same number  $n$  of equations as unknowns. However, even in that case, there is no guarantee that there is a unique solution; there can still be zero, one or many solutions (for instance, see examples in section 8.1). The case of a unique solution occurs exactly when the matrix  $A$  is non-singular.

**Theorem 9.6** *Let  $A$  be an  $n \times n$  matrix. Then*

- (i) *the homogeneous system of equations  $A\underline{\mathbf{x}} = \mathbf{0}$  has a non-zero solution if and only if  $A$  is singular;*
- (ii) *the equation system  $A\underline{\mathbf{x}} = \underline{\beta}$  has a unique solution if and only if  $A$  is non-singular.*

PROOF: (i) The solution set of the equations is exactly  $\text{nullspace}(A)$ . If  $T$  is the linear map corresponding to  $A$  then, by Corollary 5.5,

$$\text{nullspace}(T) = \ker(T) = \{\mathbf{0}\} \iff \text{nullity}(T) = 0 \iff T \text{ is non-singular,}$$

and so there are non-zero solutions if and only if  $T$  and hence  $A$  is singular.

(ii) If  $A$  is singular then its nullity is greater than 0 and so its nullspace is not equal to  $\{\mathbf{0}\}$ , and contains more than one vector. Either there are no solutions, or the solution set is  $\underline{\mathbf{x}} + \text{nullspace}(A)$  for some specific solution  $\underline{\mathbf{x}}$ , in which case there is more than one solution. Hence there cannot be a unique solution when  $A$  is singular.

Conversely, if  $A$  is non-singular, then it is invertible by Theorem 9.2, and one solution is  $\underline{\mathbf{x}} = A^{-1}\underline{\beta}$ . Since the complete solution set is then  $\underline{\mathbf{x}} + \text{nullspace}(A)$ , and  $\text{nullspace}(A) = \{\mathbf{0}\}$  in this case, the solution is unique.  $\square$

In general, it is more efficient to solve the equations  $A\underline{\mathbf{x}} = \underline{\beta}$  by elementary row operations rather than by first computing  $A^{-1}$  and then  $A^{-1}\underline{\beta}$ . However, if  $A^{-1}$  is already known for some reason, then this is a useful method.

**Example**

$$3x + 2y + z = 0 \tag{1}$$

$$4x + y + 3z = 2 \tag{2}$$

$$2x + y + 6z = 6 \tag{3}$$

Here  $A = \begin{pmatrix} 3 & 2 & 1 \\ 4 & 1 & 3 \\ 2 & 1 & 6 \end{pmatrix}$ , and we computed  $A^{-1} = \begin{pmatrix} -3/25 & 11/25 & -1/5 \\ 18/25 & -16/25 & 1/5 \\ -2/25 & -1/25 & 1/5 \end{pmatrix}$  in Section 9.

Computing  $A^{-1}\underline{\beta}$  with  $\underline{\beta} = \begin{pmatrix} 0 \\ 2 \\ 6 \end{pmatrix}$  yields the solution  $x = -8/25$ ,  $y = -2/25$ ,  $z = 28/25$ .

## 10 The Determinant of a Matrix

### 10.1 Definition of the determinant

Let  $A$  be an  $n \times n$  matrix over the field  $K$ . The *determinant* of  $A$ , which is written as  $\det(A)$  or sometimes as  $|A|$  is a certain scalar that is defined from  $A$  in a rather complicated way.

The definition for small values of  $n$  might already be familiar to you.

$$n = 1 \quad A = (\alpha) \quad \det(A) = \alpha$$

$$n = 2 \quad A = \begin{pmatrix} \alpha_{11} & \alpha_{12} \\ \alpha_{21} & \alpha_{22} \end{pmatrix} \quad \det(A) = \alpha_{11}\alpha_{22} - \alpha_{12}\alpha_{21}$$


$$n = 3 \quad A = \begin{pmatrix} \alpha_{11} & \alpha_{12} & \alpha_{13} \\ \alpha_{21} & \alpha_{22} & \alpha_{23} \\ \alpha_{31} & \alpha_{32} & \alpha_{33} \end{pmatrix}$$

$$\det(A) = \alpha_{11} \begin{vmatrix} \alpha_{22} & \alpha_{23} \\ \alpha_{32} & \alpha_{33} \end{vmatrix} - \alpha_{12} \begin{vmatrix} \alpha_{21} & \alpha_{23} \\ \alpha_{31} & \alpha_{33} \end{vmatrix} + \alpha_{13} \begin{vmatrix} \alpha_{21} & \alpha_{22} \\ \alpha_{31} & \alpha_{32} \end{vmatrix} =$$

$$\alpha_{11}\alpha_{22}\alpha_{33} - \alpha_{11}\alpha_{23}\alpha_{32} - \alpha_{12}\alpha_{21}\alpha_{33} + \alpha_{12}\alpha_{23}\alpha_{31} + \alpha_{13}\alpha_{21}\alpha_{32} - \alpha_{13}\alpha_{22}\alpha_{31}$$

The geometrical motivation for the determinant is that it represents area or volume. For  $n = 2$ , consider the position vectors of two points  $(x_1, y_1)$ ,  $(x_2, y_2)$  in the plane. Then, in the diagram below, the area of the parallelogram  $OABC$  enclosed by these two vectors is

$$r_1 r_2 \sin(\theta_2 - \theta_1) = r_1 r_2 (\sin \theta_2 \cos \theta_1 - \sin \theta_1 \cos \theta_2) = x_1 y_2 - x_2 y_1 = \begin{vmatrix} x_1 & x_2 \\ y_1 & y_2 \end{vmatrix}$$



Similarly, when  $n = 3$  the volume of the parallelepiped enclosed by the three position vectors in space is equal to (plus or minus) the determinant of the  $3 \times 3$  matrix defined by the co-ordinates of the three points.

Now we turn to the general definition for  $n \times n$  matrices. Suppose that we take the product of  $n$  entries from the matrix, where we take exactly one entry from each row and one from each column. Such a product is called an *elementary product*. There are  $n!$  such products altogether (we shall see why shortly) and the determinant is the sum of  $n!$  terms, each of which is plus or minus one of these elementary products. We say that it is a sum of  $n!$  *signed elementary products*. You should check that this holds in the 2 and 3-dimensional cases written out above.

Before we can be more precise about this, and determine which signs we choose for which elementary products, we need to make a short digression to study permutations of finite sets. A permutation of a set, which we shall take here to be the set  $X_n = \{1, 2, 3, \dots, n\}$ , is simply a bijection from  $X_n$  to itself. The set of all such permutations of  $X_n$  is called the *symmetric group*  $S_n$ . There are  $n!$  permutations altogether, so  $|S_n| = n!$ .

(A *group* is a set of objects, any two of which can be multiplied or composed together, and such that there is an identity element, and all elements have inverses. Other examples of

groups that we have met in this course are the  $n \times n$  invertible matrices over  $K$ , for any fixed  $n$ , and any field  $K$ . The study of groups, which is known as **Group Theory**, is an important branch of mathematics, but it is not the main topic of this course!

Now an elementary product contains one entry from each row of  $A$ , so let the entry in the product from the  $i$ -th row be  $\alpha_{i\phi(i)}$ . Since the product also contains exactly one entry from each column, each integer  $j \in X_n$  must occur exactly once as  $\phi(i)$ . But this is just saying that  $\phi : X_n \rightarrow X_n$  is a bijection; that is  $\phi \in S_n$ .

So an elementary product has the general form  $\alpha_{1\phi(1)}\alpha_{2\phi(2)} \cdots \alpha_{n\phi(n)}$  for some  $\phi \in S_n$ , and there are  $n!$  elementary products altogether. We want to define

$$\det(A) = \sum_{\phi \in S_n} \pm \alpha_{1\phi(1)}\alpha_{2\phi(2)} \cdots \alpha_{n\phi(n)},$$

but we still have to decide which of the elementary products has a plus sign and which has a minus sign. In fact this depends on the *sign* of the permutation  $\phi$ , which we must now define.

A *transposition* is a permutation of  $X_n$  that interchanges two numbers  $i$  and  $j$  in  $X_n$  and leaves all other numbers fixed. It is written as  $(i, j)$ . There is a theorem, which is quite easy, but we will not prove it here because it is a theorem in Group Theory, that says that every permutation can be written as a composite of transpositions. For example, if  $n = 5$ , then the permutation  $\phi$  defined by

$$\phi(1) = 4, \phi(2) = 5, \phi(3) = 3, \phi(4) = 2, \phi(5) = 1$$

is equal to the composite  $(1, 4) \circ (2, 4) \circ (2, 5)$ . (Remember that permutations are functions  $X_n \rightarrow X_n$ , so this means first apply the function  $(2, 5)$  (which interchanges 2 and 5) then apply  $(2, 4)$  and finally apply  $(1, 4)$ .)

Now a permutation  $\phi$  is said to be *even*, and to have sign  $+1$ , if  $\phi$  is a composite of an even number of permutations; and  $\phi$  is said to be *odd*, and to have sign  $-1$ , if  $\phi$  is a composite of an odd number of permutations. For example, the permutation  $\phi$  defined on  $X_n$  above is a composite of 3 transpositions, so  $\phi$  is odd and  $\text{sign}(\phi) = -1$ . The identity permutation, which leaves all points fixed, is even (because it is a composite of 0 transpositions).

Now at last we can give the general definition of the determinant.

**Definition**  $\det(A) = \sum_{\phi \in S_n} \text{sign}(\phi)\alpha_{1\phi(1)}\alpha_{2\phi(2)} \cdots \alpha_{n\phi(n)}$ .

(**Note:** You might be worrying about whether the same permutation could be both even and odd. Well, there is a moderately difficult theorem in Group Theory, which we shall not prove here, that says that this cannot happen; in other words, the concepts of even and odd permutation are *well-defined*.)

## 10.2 The effect of matrix operations on the determinant

### Theorem 10.1

- (i)  $\det(I_n) = 1$ .
- (ii) Let  $B$  result from  $A$  by applying (R2) (interchanging two rows). Then  $\det(B) = -\det(A)$ .
- (iii) If  $A$  has two equal rows then  $\det(A) = 0$ .
- (iv) Let  $B$  result from  $A$  by applying (R1) (adding a multiple of one row to another). Then  $\det(B) = \det(A)$ .
- (v) Let  $B$  result from  $A$  by applying (R3) (multiplying a row by a scalar  $\lambda$ ). Then  $\det(B) = \lambda \det(A)$ .

PROOF: (i) If  $A = I_n$  then  $\alpha_{ij} = 0$  when  $i \neq j$ . So the only non-zero elementary product in the sum occurs when  $\phi$  is the identity permutation. Hence  $\det(A) = \alpha_{11}\alpha_{22} \dots \alpha_{nn} = 1$ .

(ii) To keep the notation simple, we shall suppose that we interchange the first two rows, but the same argument works for interchanging any pair of rows. Then if  $B = (\beta_{ij})$ , we have  $\beta_{1j} = \alpha_{2j}$  and  $\beta_{2j} = \alpha_{1j}$  for all  $j$ . Hence

$$\det(B) = \sum_{\phi \in S_n} \text{sign}(\phi) \beta_{1\phi(1)} \beta_{2\phi(2)} \dots \beta_{n\phi(n)} = \sum_{\phi \in S_n} \text{sign}(\phi) \alpha_{1\phi(2)} \alpha_{2\phi(1)} \alpha_{3\phi(3)} \dots \alpha_{n\phi(n)}.$$

For  $\phi \in S_n$ , let  $\psi = \phi \circ (1, 2)$ , so  $\phi(1) = \psi(2)$  and  $\phi(2) = \psi(1)$ , and  $\text{sign}(\psi) = -\text{sign}(\phi)$ . Now, as  $\phi$  runs through all permutations in  $S_n$ , so does  $\psi$  (but in a different order), so summing over all  $\phi \in S_n$  is the same as summing over all  $\psi \in S_n$ . Hence

$$\det(B) = \sum_{\phi \in S_n} -\text{sign}(\psi) \alpha_{1\psi(1)} \alpha_{2\psi(2)} \dots \alpha_{n\psi(n)} = \sum_{\psi \in S_n} -\text{sign}(\psi) \alpha_{1\psi(1)} \alpha_{2\psi(2)} \dots \alpha_{n\psi(n)} = -\det(A).$$

(iii) Follows from (ii), because interchanging the two equal rows would give the same matrix, and hence  $\det(A) = -\det(A)$ .

(iv) Again, to simplify notation, suppose that we replace the second row  $\mathbf{r}_2$  by  $\mathbf{r}_2 + \lambda \mathbf{r}_1$  for some  $\lambda \in K$ . Then

$$\begin{aligned} \det(B) &= \sum_{\phi \in S_n} \text{sign}(\phi) \alpha_{1\phi(1)} (\alpha_{2\phi(2)} + \lambda \alpha_{1\phi(2)}) \alpha_{3\phi(3)} \dots \alpha_{n\phi(n)} \\ &= \sum_{\phi \in S_n} \text{sign}(\phi) \alpha_{1\phi(1)} \alpha_{2\phi(2)} \dots \alpha_{n\phi(n)} + \lambda \sum_{\phi \in S_n} \text{sign}(\phi) \alpha_{1\phi(1)} \alpha_{1\phi(2)} \dots \alpha_{n\phi(n)}. \end{aligned}$$

Now the first term in this sum is  $\det(A)$ , and the second is  $\lambda \det(C)$ , where  $C$  is a matrix in which the first two rows are equal. Hence  $\det(C) = 0$  by (iii), and  $\det(B) = \det(A)$ .

(v) Easy. Note that this holds even when the scalar  $\lambda = 0$ . □

**Definition** A matrix is called *upper triangular* if all of its entries below the main diagonal are zero; that is, if  $\alpha_{ij} = 0$  for all  $i > j$ .

The matrix is called *diagonal* if all entries not on the main diagonal are zero; that is,  $\alpha_{ij} = 0$  for  $i \neq j$ .

For example,  $\begin{pmatrix} 3 & 0 & -1/2 \\ 0 & -1 & -11 \\ 0 & 0 & -2/5 \end{pmatrix}$  is upper triangular, and  $\begin{pmatrix} 0 & 0 & 0 \\ 0 & 17 & 0 \\ 0 & 0 & -3 \end{pmatrix}$  is diagonal.

**Corollary 10.2** *If  $A = (\alpha_{ij})$  is upper triangular, then  $\det(A) = \alpha_{11}\alpha_{22} \dots \alpha_{nn}$  is the product of the entries on the main diagonal of  $A$ .*

PROOF: This is not hard to prove directly from the definition of the determinant. Alternatively, we can apply row operations (R1) to reduce the matrix to the diagonal matrix with the same entries  $\alpha_{ii}$  on the main diagonal, and then the result follows from parts (i) and (v) of the theorem. □

The above theorem and corollary provide the most efficient way of computing  $\det(A)$ , at least for  $n \geq 3$ . (For  $n = 2$ , it is easiest to do it straight from the definition.) Use row operations (R1) and (R2) to reduce  $A$  to upper triangular form, keeping track of changes of sign in the determinant resulting from applications of (R2), and then use Corollary 10.2.

**Example**

$$\begin{aligned}
 \begin{vmatrix} 0 & 1 & 1 & 2 \\ 1 & 2 & 1 & 1 \\ 2 & 1 & 3 & 1 \\ 1 & 2 & 4 & 2 \end{vmatrix} & \xrightarrow{\mathbf{r}_2 \leftrightarrow \mathbf{r}_1} = \begin{vmatrix} 1 & 2 & 1 & 1 \\ 0 & 1 & 1 & 2 \\ 2 & 1 & 3 & 1 \\ 1 & 2 & 4 & 2 \end{vmatrix} \xrightarrow{\substack{\mathbf{r}_3 \rightarrow \mathbf{r}_3 - 2\mathbf{r}_1 \\ \mathbf{r}_4 \rightarrow \mathbf{r}_4 - \mathbf{r}_1}} = \begin{vmatrix} 1 & 2 & 1 & 1 \\ 0 & 1 & 1 & 2 \\ 0 & -3 & 1 & -1 \\ 0 & 0 & 3 & 1 \end{vmatrix} \xrightarrow{\mathbf{r}_3 \rightarrow \mathbf{r}_3 + 3\mathbf{r}_2} = \\
 - \begin{vmatrix} 1 & 2 & 1 & 1 \\ 0 & 1 & 1 & 2 \\ 0 & 0 & 4 & 5 \\ 0 & 0 & 3 & 1 \end{vmatrix} & \xrightarrow{\mathbf{r}_4 \rightarrow \mathbf{r}_4 - 3\mathbf{r}_3/4} = \begin{vmatrix} 1 & 2 & 1 & 1 \\ 0 & 1 & 1 & 2 \\ 0 & 0 & 4 & 5 \\ 0 & 0 & 0 & -\frac{11}{4} \end{vmatrix} = 11
 \end{aligned}$$

We could have been a little more clever, and stopped the row reduction one step before the end, noticing that the determinant was equal to  $-\begin{vmatrix} 4 & 5 \\ 3 & 1 \end{vmatrix} = 11$ .

**Definition** Let  $A = (\alpha_{ij})$  be an  $m \times n$  matrix. We define the *transpose*  $A^T$  of  $A$  to be the  $n \times m$  matrix  $(\beta_{ij})$ , where  $\beta_{ij} = \alpha_{ji}$  for  $1 \leq i \leq n$ ,  $1 \leq j \leq m$ .

For example  $\begin{pmatrix} 1 & 3 & 5 \\ -2 & 0 & 6 \end{pmatrix}^T = \begin{pmatrix} 1 & -2 \\ 3 & 0 \\ 5 & 6 \end{pmatrix}$ .

**Theorem 10.3** Let  $A = (\alpha_{ij})$  be an  $n \times n$  matrix. Then  $\det(A^T) = \det(A)$ .

PROOF: Let  $A^T = (\beta_{ij})$  where  $\beta_{ij} = \alpha_{ji}$ . Then

$$\det(A^T) = \sum_{\phi \in S_n} \text{sign}(\phi) \beta_{1\phi(1)} \beta_{2\phi(2)} \cdots \beta_{n\phi(n)} = \sum_{\phi \in S_n} \text{sign}(\phi) \alpha_{\phi(1)1} \alpha_{\phi(2)2} \cdots \alpha_{\phi(n)n}.$$

Now, by rearranging the terms in the elementary product, we have

$$\alpha_{\phi(1)1} \alpha_{\phi(2)2} \cdots \alpha_{\phi(n)n} = \alpha_{1\phi^{-1}(1)} \alpha_{2\phi^{-1}(2)} \cdots \alpha_{n\phi^{-1}(n)},$$

where  $\phi^{-1}$  is the *inverse* permutation to  $\phi$ . Notice also that if  $\phi$  is a composite  $\tau_1 \circ \tau_2 \circ \cdots \circ \tau_r$  of transpositions  $\tau_i$ , then  $\phi^{-1} = \tau_r \circ \cdots \circ \tau_2 \circ \tau_1$  (because each  $\tau_i \circ \tau_i$  is the identity permutation). Hence  $\text{sign}(\phi) = \text{sign}(\phi^{-1})$ . Also, summing over all  $\phi \in S_n$  is the same as summing over all  $\phi^{-1} \in S_n$ , so we have

$$\begin{aligned}
 \det(A^T) &= \sum_{\phi \in S_n} \text{sign}(\phi) \alpha_{1\phi^{-1}(1)} \alpha_{2\phi^{-1}(2)} \cdots \alpha_{n\phi^{-1}(n)} \\
 &= \sum_{\phi^{-1} \in S_n} \text{sign}(\phi^{-1}) \alpha_{1\phi^{-1}(1)} \alpha_{2\phi^{-1}(2)} \cdots \alpha_{n\phi^{-1}(n)} = \det(A).
 \end{aligned}$$

□

If you find proofs like the above, where we manipulate sums of products, hard to follow, then it might be helpful to write it out in full in a small case, such as  $n = 3$ . Then

$$\begin{aligned}
 \det(A^T) &= \beta_{11}\beta_{22}\beta_{33} - \beta_{11}\beta_{23}\beta_{32} - \beta_{12}\beta_{21}\beta_{33} + \beta_{12}\beta_{23}\beta_{31} + \beta_{13}\beta_{21}\beta_{32} - \beta_{13}\beta_{22}\beta_{31} \\
 &= \alpha_{11}\alpha_{22}\alpha_{33} - \alpha_{11}\alpha_{32}\alpha_{23} - \alpha_{21}\alpha_{12}\alpha_{33} + \alpha_{21}\alpha_{32}\alpha_{13} + \alpha_{31}\alpha_{12}\alpha_{23} - \alpha_{31}\alpha_{22}\alpha_{13} \\
 &= \alpha_{11}\alpha_{22}\alpha_{33} - \alpha_{11}\alpha_{23}\alpha_{32} - \alpha_{12}\alpha_{21}\alpha_{33} + \alpha_{12}\alpha_{23}\alpha_{31} + \alpha_{13}\alpha_{21}\alpha_{32} - \alpha_{13}\alpha_{22}\alpha_{31} \\
 &= \det(A).
 \end{aligned}$$

**Corollary 10.4** All of Theorem 10.1 remains true if we replace rows by columns.

PROOF: This follows from Theorems 10.1 and 10.3, because we can apply column operations to  $A$  by transposing it, applying the corresponding row operations, and then re-transposing it.  $\square$

**Theorem 10.5** For an  $n \times n$  matrix  $A$ ,  $\det(A) = 0$  if and only if  $A$  is singular.

PROOF:  $A$  can be reduced to row reduced echelon form by using row operations (R1), (R2) and (R3). By Theorem 8.5, none of these operations affect the rank of  $A$ , and so they do not affect whether or not  $A$  is singular (remember ‘singular’ means  $\text{rank}(A) < n$ ; see definition after Corollary 5.5). By Theorem 10.1, they do not affect whether or not  $\det(A) = 0$ . So we can assume that  $A$  is in row reduced echelon form.

Then  $\text{rank}(A)$  is the number of non-zero rows of  $A$ , so if  $A$  is singular then it has some zero rows. But then  $\det(A) = 0$ . On the other hand, if  $A$  is nonsingular then, as we saw in Section 9.2, the fact that  $A$  is in row reduced echelon form implies that  $A = I_n$ , so  $\det(A) = 1 \neq 0$ .  $\square$

### 10.3 The determinant of a product

**Example** Let  $A = \begin{pmatrix} 1 & 2 \\ 3 & 2 \end{pmatrix}$  and  $B = \begin{pmatrix} -1 & -1 \\ 2 & 0 \end{pmatrix}$ . Then  $\det(A) = -4$  and  $\det(B) = 2$ .

We have  $A + B = \begin{pmatrix} 0 & 1 \\ 5 & 2 \end{pmatrix}$  and  $\det(A + B) = -5 \neq \det(A) + \det(B)$ . In fact, in general there is no simple relationship between  $\det(A + B)$  and  $\det(A)$ ,  $\det(B)$ .

However,  $AB = \begin{pmatrix} 3 & -1 \\ 1 & -3 \end{pmatrix}$ , and  $\det(AB) = -8 = \det(A)\det(B)$ .

In this subsection, we shall prove that this simple relationship holds in general.

Recall from Section 9.3 the definition of an *elementary* matrix  $E$ , and the property that if we multiply a matrix  $B$  on the left by  $E$ , then the effect is to apply the corresponding elementary row operation to  $B$ . This enables us to prove:

**Lemma 10.6** If  $E$  is an  $n \times n$  elementary matrix, and  $B$  is any  $n \times n$  matrix, then  $\det(EB) = \det(E)\det(B)$ .

PROOF:  $E$  is one of the three types  $E(n)_{\lambda,ij}^1$ ,  $E(n)_{ij}^2$  or  $E(n)_{\lambda,i}^3$ , and multiplying  $B$  on the left by  $E$  has the effect of applying (R1), (R2) or (R3) to  $B$ , respectively. Hence, by Theorem 10.1,  $\det(EB) = \det(B)$ ,  $-\det(B)$ , or  $\lambda\det(B)$ , respectively. But by considering the special case  $B = I_n$ , we see that  $\det(E) = 1, -1$  or  $\lambda$ , respectively, and so  $\det(EB) = \det(E)\det(B)$  in all three cases.  $\square$

**Theorem 10.7** For any two  $n \times n$  matrices  $A$  and  $B$ , we have  $\det(AB) = \det(A)\det(B)$ .

PROOF: Suppose first that  $\det(A) = 0$ . Then we have  $\text{rank}(A) < n$  by Theorem 10.5. Let  $T_1, T_2 : V \rightarrow V$  be linear maps corresponding to  $A$  and  $B$ , where  $\dim(V) = n$ . Then  $AB$  corresponds to  $T_1T_2$  (by Theorem 7.3). By Corollary 5.5,  $\text{rank}(A) = \text{rank}(T_1) < n$  implies that  $T_1$  is not surjective. But then  $T_1T_2$  cannot be surjective, so  $\text{rank}(T_1T_2) = \text{rank}(AB) < n$ . Hence  $\det(AB) = 0$  so  $\det(AB) = \det(A)\det(B)$ .

On the other hand, if  $\det(A) \neq 0$ , then  $A$  is nonsingular, and hence invertible, so by Theorem 9.5  $A$  is a product  $E_1E_2 \dots E_r$  of elementary matrices  $E_i$ . Hence  $\det(AB) = \det(E_1E_2 \dots E_rB)$ . Now the result follows from the above lemma, because

$$\begin{aligned} \det(AB) &= \det(E_1)\det(E_2 \dots E_rB) = \det(E_1)\det(E_2)\det(E_3 \dots E_rB) = \\ &= \det(E_1)\det(E_2) \dots \det(E_r)\det(B) = \det(E_1E_2 \dots E_r)\det(B) = \det(A)\det(B). \end{aligned} \quad \square$$

## 10.4 Minors and cofactors

Let  $A = (\alpha_{ij})$  be an  $n \times n$  matrix. Let  $A_{ij}$  be the  $(n-1) \times (n-1)$  matrix obtained from  $A$  by deleting the  $i$ -th row and the  $j$ -th column of  $A$ . Now let  $M_{ij} = \det(A_{ij})$ . Then  $M_{ij}$  is called the  $(i, j)$ -th *minor* of  $A$ .

For example, if  $A = \begin{pmatrix} 2 & 1 & 0 \\ 3 & -1 & 2 \\ 5 & -2 & 0 \end{pmatrix}$ , then  $A_{12} = \begin{pmatrix} 3 & 2 \\ 5 & 0 \end{pmatrix}$  and  $A_{31} = \begin{pmatrix} 1 & 0 \\ -1 & 2 \end{pmatrix}$ , and so  $M_{12} = -10$  and  $M_{31} = 2$ .

We define  $c_{ij}$  to be equal to  $M_{ij}$  if  $i + j$  is even, and to  $-M_{ij}$  if  $i + j$  is odd. Or, more concisely,

$$c_{ij} = (-1)^{i+j} M_{ij} = (-1)^{i+j} \det(A_{ij}).$$

Then  $c_{ij}$  is called the  $(i, j)$ -th *cofactor* of  $A$ . In the example above,

$$\begin{aligned} c_{11} &= \begin{vmatrix} -1 & 2 \\ -2 & 0 \end{vmatrix} = 4, & c_{12} &= -\begin{vmatrix} 3 & 2 \\ 5 & 0 \end{vmatrix} = 10, & c_{13} &= \begin{vmatrix} 3 & -1 \\ 5 & -2 \end{vmatrix} = -1 \\ c_{21} &= -\begin{vmatrix} 1 & 0 \\ -2 & 0 \end{vmatrix} = 0, & c_{22} &= \begin{vmatrix} 2 & 0 \\ 5 & 0 \end{vmatrix} = 0, & c_{23} &= -\begin{vmatrix} 2 & 1 \\ 5 & -2 \end{vmatrix} = 9 \\ c_{31} &= \begin{vmatrix} 1 & 0 \\ -1 & 2 \end{vmatrix} = 2, & c_{32} &= -\begin{vmatrix} 2 & 0 \\ 3 & 2 \end{vmatrix} = -4, & c_{33} &= \begin{vmatrix} 2 & 1 \\ 3 & -1 \end{vmatrix} = -5. \end{aligned}$$

**Theorem 10.8** *Let  $A$  be an  $n \times n$  matrix.*

(i) *(Expansion of a determinant by the  $i$ -th row.) For any  $i$  with  $1 \leq i \leq n$ , we have*

$$\det(A) = \alpha_{i1}c_{i1} + \alpha_{i2}c_{i2} + \cdots + \alpha_{in}c_{in} = \sum_{j=1}^n \alpha_{ij}c_{ij}.$$

(ii) *(Expansion of a determinant by the  $j$ -th column.) For any  $j$  with  $1 \leq j \leq n$ , we have*

$$\det(A) = \alpha_{1j}c_{1j} + \alpha_{2j}c_{2j} + \cdots + \alpha_{nj}c_{nj} = \sum_{i=1}^n \alpha_{ij}c_{ij}.$$

For example, expanding the determinant of the matrix  $A$  above by the first row, the third row, and the second column give respectively:

$$\begin{aligned} \det(A) &= 2 \times 4 + 1 \times 10 + 0 \times -1 = 18 \\ \det(A) &= 5 \times 2 + -2 \times -4 + 0 \times -5 = 18 \\ \det(A) &= 1 \times 10 + -1 \times 0 + -2 \times -4 = 18 \end{aligned}$$

PROOF OF THEOREM 10.8: By definition, we have

$$\det(A) = \sum_{\phi \in S_n} \text{sign}(\phi) \alpha_{1\phi(1)} \alpha_{2\phi(2)} \cdots \alpha_{n\phi(n)} \quad (*)$$

**Step 1.** We first find the sum of all of those signed elementary products in the sum (\*) that contain  $\alpha_{nn}$ . These arise from those permutations  $\phi$  with  $\phi(n) = n$ ; so the required sum is

$$\begin{aligned} \sum_{\substack{\phi \in S_n \\ \phi(n) = n}} \text{sign}(\phi) \alpha_{1\phi(1)} \alpha_{2\phi(2)} \cdots \alpha_{n\phi(n)} &= \alpha_{nn} \sum_{\phi \in S_{n-1}} \text{sign}(\phi) \alpha_{1\phi(1)} \alpha_{2\phi(2)} \cdots \alpha_{n-1\phi(n-1)} \\ &= \alpha_{nn} M_{nn} = \alpha_{nn} c_{nn}. \end{aligned}$$

**Step 2.** Next we fix any  $i$  and  $j$  with  $1 \leq i, j \leq n$ , and find the sum of all of those signed elementary products in the sum (\*) that contain  $\alpha_{ij}$ . We move row  $\mathbf{r}_i$  of  $A$  to  $\mathbf{r}_n$  by interchanging  $\mathbf{r}_i$  with  $\mathbf{r}_{i+1}, \mathbf{r}_{i+2}, \dots, \mathbf{r}_n$  in turn. This involves  $n - i$  applications of (R2), and leaves the rows of  $A$  other than  $\mathbf{r}_i$  in their original order. We then move column  $\mathbf{c}_j$  to  $\mathbf{c}_n$  in the same way, by applying (C2)  $n - j$  times. Let the resulting matrix be  $B = (\beta_{ij})$  and denote its minors by  $N_{ij}$ . Then  $\beta_{nn} = \alpha_{ij}$ , and  $N_{nn} = M_{ij}$ . Furthermore,

$$\det(B) = (-1)^{2n-i-j} \det(A) = (-1)^{i+j} \det(A),$$

because  $(2n - i - j) - (i + j)$  is even.

Now, by the result of Step 1, the sum of terms in  $\det(B)$  involving  $\beta_{nn}$  is

$$\beta_{nn} N_{nn} = \alpha_{ij} M_{ij} = (-1)^{i+j} \alpha_{ij} c_{ij},$$

and hence, since  $\det(B) = (-1)^{i+j} \det(A)$ , the sum of terms involving  $\alpha_{ij}$  in  $\det(A)$  is  $\alpha_{ij} c_{ij}$ .

**Step 3.** The result follows from Step 2, because every signed elementary product in the sum (\*) involves exactly one array element  $\alpha_{ij}$  from each row and from each column. Hence, for any given row or column, we get the full sum (\*) by adding up the total of those products involving each individual element in that row or column.  $\square$

Expanding by a row and column can sometimes be a quick method of evaluating the determinant of matrices containing a lot of zeros. For example, let

$$A = \begin{pmatrix} 9 & 0 & 2 & 6 \\ 1 & 2 & 9 & -3 \\ 0 & 0 & -2 & 0 \\ -1 & 0 & -5 & 2 \end{pmatrix}.$$

Then, expanding by the third row, we get  $\det(A) = -2 \begin{vmatrix} 9 & 0 & 6 \\ 1 & 2 & -3 \\ -1 & 0 & 2 \end{vmatrix}$ , and then expanding

by the second column,  $\det(A) = -2 \times 2 \begin{vmatrix} 9 & 6 \\ -1 & 2 \end{vmatrix} = -96$ .

## 10.5 The inverse of a matrix using determinants

**Definition** Let  $A$  be an  $n \times n$  matrix. We define the *adjoint* matrix  $\text{adj}(A)$  of  $A$  to be the  $n \times n$  matrix of which the  $(i, j)$ -th element is the cofactor  $c_{ji}$ . In other words, it is the transpose of the matrix of cofactors.

In the example above,

$$A = \begin{pmatrix} 2 & 1 & 0 \\ 3 & -1 & 2 \\ 5 & -2 & 0 \end{pmatrix}, \quad \text{adj}(A) = \begin{pmatrix} 4 & 0 & 2 \\ 10 & 0 & -4 \\ -1 & 9 & -5 \end{pmatrix}.$$

**Theorem 10.9**  $A \text{adj}(A) = \det(A) I_n = \text{adj}(A) A$

PROOF: Let  $B = A \text{adj}(A) = (\beta_{ij})$ . Then  $\beta_{ii} = \sum_{k=1}^n \alpha_{ik} c_{ik} = \det(A)$  by Theorem 10.8 (expansion by the  $i$ -th row of  $A$ ). For  $i \neq j$ ,  $\beta_{ij} = \sum_{k=1}^n \alpha_{ik} c_{jk}$ , which is the determinant of a matrix  $C$  obtained from  $A$  by substituting the  $i$ -th row of  $A$  for the  $j$ -th row. But then  $C$  has two equal rows, so  $\beta_{ij} = \det(C) = 0$  by Theorem 10.1(iii). (This is sometimes called an expansion by *alien cofactors*.) Hence  $A \text{adj}(A) = \det(A) I_n$ . A similar argument using columns instead of rows gives  $\text{adj}(A) A = \det(A) I_n$ .  $\square$

In the example above, check that  $A \text{adj}(A) = \text{adj}(A) A = 18I_3$ .

**Corollary 10.10** *If  $\det(A) \neq 0$ , then  $A^{-1} = \frac{1}{\det(A)}\text{adj}(A)$ .*

(Theorems 9.2 and 10.5 imply that  $A$  is invertible if and only if  $\det(A) \neq 0$ .)

So, in the example above,

$$\begin{pmatrix} 2 & 1 & 0 \\ 3 & -1 & 2 \\ 5 & -2 & 0 \end{pmatrix}^{-1} = \frac{1}{18} \begin{pmatrix} 4 & 0 & 2 \\ 10 & 0 & -4 \\ -1 & 9 & -5 \end{pmatrix},$$

and in the example in Section 9,

$$A = \begin{pmatrix} 3 & 2 & 1 \\ 4 & 1 & 3 \\ 2 & 1 & 6 \end{pmatrix}, \quad \text{adj}(A) = \begin{pmatrix} 3 & -11 & 5 \\ -18 & 16 & -5 \\ 2 & 1 & -5 \end{pmatrix}, \quad \det(A) = -25, \quad A^{-1} = \frac{-1}{25}\text{adj}(A).$$

For  $2 \times 2$  and (possibly)  $3 \times 3$  matrices, the cofactor method of computing the inverse is often the quickest. For larger matrices, the row reduction method described in Section 9 is quicker.

## 10.6 Cramer's rule for solving simultaneous equations

Given a system  $A\mathbf{x} = \underline{\beta}$  of  $n$  equations in  $n$  unknowns, where  $A = (\alpha_{ij})$  is non-singular, the solution is  $\mathbf{x} = A^{-1}\underline{\beta}$ . So the  $i$ -th component  $x_i$  of this column vector is the  $i$ -th row of  $A^{-1}\underline{\beta}$ . Now, by Corollary 10.10,  $A^{-1} = \frac{1}{\det(A)}\text{adj}(A)$ , and its  $(i, j)$ -th entry is  $c_{ji}/\det(A)$ . Hence

$$x_i = \frac{1}{\det(A)} \sum_{j=1}^n c_{ji}\beta_j.$$

Now let  $A_i$  be the matrix obtained from  $A$  by substituting  $\underline{\beta}$  for the  $i$ -th column of  $A$ . Then the sum  $\sum_{j=1}^n c_{ji}\beta_j$  is precisely the expansion of  $\det(A_i)$  by its  $i$ -th column (see Theorem 10.8). Hence we have  $x_i = \det(A_i)/\det(A)$ . This is Cramer's rule.

This is more of a curiosity than a practical method of solving simultaneous equations, although it can be quite quick in the  $2 \times 2$  case. Even in the  $3 \times 3$  case it is rather slow.

**Example**

$$\begin{array}{rcl} 2x & + & z = 1 \\ & y & - 2z = 0 \\ x & + & y + z = -1 \end{array}$$

$$\det(A) = \begin{vmatrix} 2 & 0 & 1 \\ 0 & 1 & -2 \\ 1 & 1 & 1 \end{vmatrix} = 5, \quad \det(A_1) = \begin{vmatrix} 1 & 0 & 1 \\ 0 & 1 & -2 \\ -1 & 1 & 1 \end{vmatrix} = 4$$

$$\det(A_2) = \begin{vmatrix} 2 & 1 & 1 \\ 0 & 0 & -2 \\ 1 & -1 & 1 \end{vmatrix} = -6, \quad \det(A_3) = \begin{vmatrix} 2 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & -1 \end{vmatrix} = -3$$

so the solution is  $x = 4/5$ ,  $y = -6/5$ ,  $z = -3/5$ .

## 11 Change of Basis and Equivalent Matrices

In this section, we investigate the relationship between the matrices corresponding to the same linear map  $T : U \rightarrow V$ , but using different bases for the vector spaces  $U$  and  $V$ . We first discuss the relation between two different bases of the same space. Assume throughout the section that all vector spaces are over the same field  $K$ .

Let  $U$  be a vector space of dimension  $n$ , and let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  and  $\mathbf{e}'_1, \dots, \mathbf{e}'_n$  be two bases of  $U$ . The matrix  $P$  of the identity map  $I_U : U \rightarrow U$  using the basis  $\mathbf{e}_1, \dots, \mathbf{e}_n$  in the domain and  $\mathbf{e}'_1, \dots, \mathbf{e}'_n$  in the range is called the *change of basis matrix* from the basis of  $\mathbf{e}_i$ -s to the basis of  $\mathbf{e}'_i$ -s.

Let us look carefully what this definition says. Taking  $P = (\sigma_{ij})$ , we obtain from Section 7.1

$$I_U(\mathbf{e}_j) = \mathbf{e}_j = \sum_{i=1}^n \sigma_{ij} \mathbf{e}'_i \quad \text{for } 1 \leq j \leq n. \quad (*)$$

In other words, the columns of  $P$  are the basis vectors  $\mathbf{e}_i$  as column vectors in the  $\mathbf{e}'_i$ .

**Proposition 11.1** *The change of basis matrix is invertible. More precisely, if  $P$  is the change of basis matrix from the basis of  $\mathbf{e}_i$ -s to the basis of  $\mathbf{e}'_i$ -s and  $Q$  is the change of basis matrix from the basis of  $\mathbf{e}'_i$ -s to the basis of  $\mathbf{e}_i$ -s then  $P = Q^{-1}$ .*

PROOF: Consider the composition  $I_U : U \xrightarrow{I_U} U \xrightarrow{I_U} U$  using the basis of  $\mathbf{e}'_i$ -s for the first and the third copy of  $U$  and the basis of  $\mathbf{e}_i$ -s for the middle copy of  $U$ . The composition has matrix  $I_n$  because the same basis is used for both domain and range. But the first  $I_U$  has matrix  $Q$  (change of basis from  $\mathbf{e}'_i$ -s to  $\mathbf{e}_i$ -s) and the second  $I_U$  similarly has matrix  $P$ . Therefore by Theorem 7.3,  $I_n = PQ$ .

Similarly,  $I_n = QP$ . Consequently,  $P = Q^{-1}$ .  $\square$

**Example** Let  $U = \mathbb{R}^3$ ,  $\mathbf{e}'_1 = (1, 0, 0)$ ,  $\mathbf{e}'_2 = (0, 1, 0)$ ,  $\mathbf{e}'_3 = (0, 0, 1)$  (the standard basis) and  $\mathbf{e}_1 = (0, 2, 1)$ ,  $\mathbf{e}_2 = (1, 1, 0)$ ,  $\mathbf{e}_3 = (1, 0, 0)$ . Then

$$P = \begin{pmatrix} 0 & 1 & 1 \\ 2 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

**Proposition 11.2** *With the above notation, let  $\mathbf{v} \in U$ , and let  $\underline{\mathbf{v}}$  and  $\underline{\mathbf{v}}'$  denote the column vectors associated with  $\mathbf{v}$  when we use the bases  $\mathbf{e}_1, \dots, \mathbf{e}_n$  and  $\mathbf{e}'_1, \dots, \mathbf{e}'_n$ , respectively. Then  $P\underline{\mathbf{v}} = \underline{\mathbf{v}}'$ .*

PROOF: This follows immediately from Proposition 7.2 applied to the identity map  $I_U$ .  $\square$

Now let  $T : U \rightarrow V$  be a linear map, where  $\dim(U) = n$ ,  $\dim(V) = m$ . Choose a basis  $\mathbf{e}_1, \dots, \mathbf{e}_n$  of  $U$  and a basis  $\mathbf{f}_1, \dots, \mathbf{f}_m$  of  $V$ . Then, from Section 7.1, we have

$$T(\mathbf{e}_j) = \sum_{i=1}^m \alpha_{ij} \mathbf{f}_i \quad \text{for } 1 \leq j \leq n.$$


where  $A = (\alpha_{ij})$  is the  $m \times n$  matrix of  $T$  with respect to the bases  $\{\mathbf{e}_i\}$  and  $\{\mathbf{f}_i\}$  of  $U$  and  $V$ .

Now choose new bases  $\mathbf{e}'_1, \dots, \mathbf{e}'_n$  of  $U$  and  $\mathbf{f}'_1, \dots, \mathbf{f}'_m$  of  $V$ . Then

$$T(\mathbf{e}'_j) = \sum_{i=1}^m \beta_{ij} \mathbf{f}'_i \quad \text{for } 1 \leq j \leq n,$$

where  $B = (\beta_{ij})$  is the  $m \times n$  matrix of  $T$  with respect to the bases  $\{\mathbf{e}'_i\}$  and  $\{\mathbf{f}'_i\}$  of  $U$  and  $V$ . Our objective is to find the relationship between  $A$  and  $B$  in terms of the change of basis matrices.

Let the  $n \times n$  matrix  $P = (\sigma_{ij})$  be the change of basis matrix from  $\{\mathbf{e}_i\}$  to  $\{\mathbf{e}'_i\}$ , and let the  $m \times m$  matrix  $Q = (\tau_{ij})$  be the change of basis matrix from  $\{\mathbf{f}_i\}$  to  $\{\mathbf{f}'_i\}$ .



**Theorem 11.3** *With the above notation, we have  $BP = QA$ , or equivalently  $B = QAP^{-1}$ .*

PROOF: By Theorem 7.3,  $BP$  represents the composite of the linear maps  $I_U$  using bases  $\{\mathbf{e}_i\}$  and  $\{\mathbf{e}'_i\}$  and  $T$  using bases  $\{\mathbf{e}'_i\}$  and  $\{\mathbf{f}'_i\}$ . So  $BP$  represents  $T$  using bases  $\{\mathbf{e}_i\}$  and  $\{\mathbf{f}'_i\}$ . Similarly,  $QA$  represents the composite of  $T$  using bases  $\{\mathbf{e}_i\}$  and  $\{\mathbf{f}_i\}$  and  $I_V$  using bases  $\{\mathbf{f}_i\}$  and  $\{\mathbf{f}'_i\}$ , so  $QA$  also represents  $T$  using bases  $\{\mathbf{e}_i\}$  and  $\{\mathbf{f}'_i\}$ . Hence  $BP = QA$ .  $\square$

**Corollary 11.4** *Two  $m \times n$  matrices  $A$  and  $B$  represent the same linear map from an  $n$ -dimensional vector space to an  $m$ -dimensional vector space (with respect to different bases) if and only if there exist invertible  $n \times n$  and  $m \times m$  matrices  $P$  and  $Q$  with  $B = QAP$ .*

PROOF: It follows from the Theorem 11.3 that  $A$  and  $B$  represent the same linear map if there exist change of basis matrices  $P$  and  $Q$  with  $B = QAP^{-1}$ , and by Proposition 11.1 the change of basis matrices are precisely invertible matrices of the correct size. By replacing  $P$  by  $P^{-1}$ , we see that this is equivalent to saying that there exist invertible  $Q, P$  with  $B = QAP$ .  $\square$

**Definition** Two  $m \times n$  matrices  $A$  and  $B$  are said to be *equivalent* if there exist invertible  $P$  and  $Q$  with  $B = QAP$ ; that is, if they represent the same linear map.

It is easy to check that being equivalent is an equivalence relation on the set  $K^{m,n}$  of  $m \times n$  matrices over  $K$ . We shall show now that equivalence of matrices has other characterisations.

**Theorem 11.5** *Let  $A$  and  $B$  be  $m \times n$  matrices over  $K$ . Then the following conditions on  $A$  and  $B$  are equivalent.*

- (i)  $A$  and  $B$  are equivalent.
- (ii)  $A$  and  $B$  represent the same linear map with respect to different bases.
- (iii)  $A$  and  $B$  have the same rank.
- (iv)  $B$  can be obtained from  $A$  by application of elementary row and column operations.

PROOF: (i)  $\Leftrightarrow$  (ii): This is true by Corollary 11.4.

(ii)  $\Rightarrow$  (iii): Since  $A$  and  $B$  both represent the same linear map  $T$ , we have  $\text{rank}(A) = \text{rank}(B) = \text{rank}(T)$ .

(iii)  $\Rightarrow$  (iv): By Theorem 8.2, if  $A$  and  $B$  both have rank  $s$ , then they can both be brought into the form

$$E_s = \left( \begin{array}{c|c} I_s & \mathbf{0}_{s,n-s} \\ \hline \mathbf{0}_{m-s,s} & \mathbf{0}_{m-s,n-s} \end{array} \right)$$

by elementary row and column operations. Since these operations are invertible, we can first transform  $A$  to  $E_s$  and then transform  $E_s$  to  $B$ .

(iv)  $\Rightarrow$  (i): We saw in Section 9.2 that applying an elementary row operation to  $A$  can be achieved by multiplying  $A$  on the left by an elementary row matrix, and similarly applying an elementary column operation can be done by multiplying  $A$  on the right by an elementary column matrix. Hence (iv) implies that there exist elementary row matrices  $R_1, \dots, R_r$  and

elementary column matrices  $C_1, \dots, C_s$  with  $B = R_r \dots R_1 A C_1 \dots C_s$ . Since elementary matrices are invertible,  $Q = R_r \dots R_1$  and  $P = C_1 \dots C_s$  are invertible and  $B = QAP$ .  $\square$

In the above proof, we also showed the following:

**Proposition 11.6** *Any  $m \times n$  matrix is equivalent to the matrix  $E_s$  defined above, where  $s = \text{rank}(A)$ .*

The form  $E_s$  is known as a *canonical form* for  $m \times n$  matrices under equivalence. This means that it is an easily recognizable representative of its equivalence class.

## 11\* An Alternative Approach.

In this section we provide an alternative presentation for some of the material of Section 11. This is naturally the same content, so it can be skipped at first by the reader comfortable with the material of the previous section. We introduce here another system of notation which in turn provides a new perspective to the change of basis formula.

Again, let  $U$  and  $V$  be vector spaces over a field  $K$ , and let  $T : U \rightarrow V$  be a linear map. Suppose further that  $\mathcal{E} = \{e_1, e_2, \dots, e_n\}$  and  $\mathcal{F} = \{f_1, f_2, \dots, f_m\}$  are the bases of  $U$  and  $V$  respectively. If  $u \in U$  is equal to  $\alpha_1 e_1 + \dots + \alpha_n e_n$ , then we write

$$[u]_{\mathcal{E}} = \begin{bmatrix} \alpha_1 \\ \cdots \\ \alpha_n \end{bmatrix}.$$

That is,  $[u]_{\mathcal{E}}$  is *the expression of  $u$  with respect to the basis  $\mathcal{E}$  as a column vector*.

Let us denote a matrix of  $T$  with respect to the bases  $\mathcal{E}$  and  $\mathcal{F}$  by

$$[T]_{\mathcal{F}}^{\mathcal{E}}.$$

By definition,

$$[T]_{\mathcal{F}}^{\mathcal{E}} = [ [T(e_1)]_{\mathcal{F}} \dots [T(e_n)]_{\mathcal{F}} ]$$

-its columns are the expressions of the vectors  $T(e_1), \dots, T(e_n)$  with respect to the basis  $\mathcal{F}$ . Multiplication of matrices is defined so that multiplication by  $[T]_{\mathcal{F}}^{\mathcal{E}}$  converts the expression for a vector  $u$  with respect to the basis  $\mathcal{E}$ , into the expression for  $T(u)$  with respect to the basis  $\mathcal{F}$ . That is,

$$[T(u)]_{\mathcal{F}} = [T]_{\mathcal{F}}^{\mathcal{E}} [u]_{\mathcal{E}}.$$

Once you know the rule for multiplying a column vector by a matrix, this last equality is actually obvious:

$$\begin{aligned} [T]_{\mathcal{F}}^{\mathcal{E}} \begin{bmatrix} \alpha_1 \\ \cdots \\ \alpha_n \end{bmatrix} &= \alpha_1 \cdot \text{column 1 of } [T]_{\mathcal{F}}^{\mathcal{E}} + \dots + \alpha_n \cdot \text{column } n \text{ of } [T]_{\mathcal{F}}^{\mathcal{E}} = \\ &= \alpha_1 [T(e_1)]_{\mathcal{F}} + \dots + \alpha_n [T(e_n)]_{\mathcal{F}} = \\ &= [\alpha_1 T(e_1) + \dots + \alpha_n T(e_n)]_{\mathcal{F}} = \end{aligned}$$

$$= [T(\alpha_1 e_1 + \dots + \alpha_n e_n)]_{\mathcal{F}} = [T(u)]_{\mathcal{F}}$$

This corresponds to the Proposition 7.2 that we saw earlier. From this we are going to deduce the rest. But before we proceed with the change of basis formula, let us recall that in Section 7.2 we also proved that the matrix of a composition of linear maps equals to the product of the corresponding matrices. Let us begin by “rederiving” this fact in our new settings.

As in Theorem 7.3, let  $T_2 : U \rightarrow V$  and  $T_1 : V \rightarrow W$  be linear maps where  $U$ ,  $V$  and  $W$  are vector spaces over a field  $K$ . Suppose further that  $\mathcal{E}$  is a basis for  $U$ ,  $\mathcal{F}$  is a basis for  $V$  and  $\mathcal{G}$  is a basis for  $W$ . Then

$$[T_1]_{\mathcal{G}}^{\mathcal{F}} [T_2]_{\mathcal{F}}^{\mathcal{E}} = [T_1]_{\mathcal{G}}^{\mathcal{F}} [ [T_2(e_1)]_{\mathcal{F}} \dots [T_2(e_n)]_{\mathcal{F}} ].$$

Since for any  $v \in V$  we have

$$[T_1]_{\mathcal{G}}^{\mathcal{F}} [v]_{\mathcal{F}} = [T_1(v)]_{\mathcal{G}},$$

in particular, we obtain that

$$[T_1]_{\mathcal{G}}^{\mathcal{F}} [T_2(e_i)]_{\mathcal{F}} = [T_1(T_2(e_i))]_{\mathcal{G}}$$

Therefore

$$[T_1]_{\mathcal{G}}^{\mathcal{F}} [T_2]_{\mathcal{F}}^{\mathcal{E}} = [ [T_1(T_2(e_1))]_{\mathcal{G}} \dots [T_1(T_2(e_n))]_{\mathcal{G}} ] = [T_1 \circ T_2]_{\mathcal{G}}^{\mathcal{E}}$$

which corresponds to Theorem 7.3 which we saw earlier.

Finally, let us deal with the change of basis procedure. Suppose that  $\mathcal{E}$  and  $\mathcal{E}'$  are both bases of the same space  $U$ . Denote the identity map from  $U$  to  $U$  by  $I_U$ . Then of course,  $[I_U]_{\mathcal{E}}^{\mathcal{E}}$  and  $[I_U]_{\mathcal{E}'}^{\mathcal{E}'}$  are both identity matrices (matrices with 1's down the diagonal and zero everywhere else). For example,

$$\begin{aligned} [I_U]_{\mathcal{E}}^{\mathcal{E}} &= [ [I_U(e_1)]_{\mathcal{E}} \dots [I_U(e_n)]_{\mathcal{E}} ] = [ [e_1]_{\mathcal{E}} \dots [e_n]_{\mathcal{E}} ] = \\ &= \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & 1 \end{bmatrix} \end{aligned}$$

On the other hand,  $[I_U]_{\mathcal{E}'}^{\mathcal{E}}$  and  $[I_U]_{\mathcal{E}}^{\mathcal{E}'}$  are not identity matrices (provided,  $\mathcal{E} \neq \mathcal{E}'$ , of course). For example,

$$[I_U]_{\mathcal{E}'}^{\mathcal{E}} = [ [e_1]_{\mathcal{E}'} \dots [e_n]_{\mathcal{E}'} ]$$

and of course,  $[e_i]_{\mathcal{E}'}$  is not the  $i$ -th column of the identity matrix, unless  $e_i = e'_i$ .

However,  $[I_U]_{\mathcal{E}'}^{\mathcal{E}}$  and  $[I_U]_{\mathcal{E}}^{\mathcal{E}'}$  are mutually inverse, as

$$[I_U]_{\mathcal{E}'}^{\mathcal{E}} [I_U]_{\mathcal{E}}^{\mathcal{E}'} = [I_U \circ I_U]_{\mathcal{E}'}^{\mathcal{E}'} = [I_U]_{\mathcal{E}'}^{\mathcal{E}'}$$

is the identity matrix (please, note that this corresponds to Proposition 11.1).

Similarly, if  $\mathcal{F}$  and  $\mathcal{F}'$  are both bases of a vector space  $V$ , and  $I_V$  is the identity map on  $V$ , then  $[I_V]_{\mathcal{F}'}^{\mathcal{F}}$  and  $[I_V]_{\mathcal{F}}^{\mathcal{F}'}$  are mutually inverse.

Finally, let  $T : U \rightarrow V$  be a linear map. Then obviously,

$$T \circ I_U = I_V \circ T.$$

Hence,

$$[T \circ I_U]_{\mathcal{F}'}^{\mathcal{E}} = [I_V \circ T]_{\mathcal{F}'}^{\mathcal{E}}$$

and so

$$[T]_{\mathcal{F}'}^{\mathcal{E}'} [I_U]_{\mathcal{E}'}^{\mathcal{E}} = [I_V]_{\mathcal{F}'}^{\mathcal{F}} [T]_{\mathcal{F}}^{\mathcal{E}}.$$

Multiplying both sides on the right by the inverse of  $[I_U]_{\mathcal{E}'}^{\mathcal{E}}$ , we obtain

$$[T]_{\mathcal{F}'}^{\mathcal{E}'} = [I_V]_{\mathcal{F}'}^{\mathcal{F}} [T]_{\mathcal{F}}^{\mathcal{E}} [I_U]_{\mathcal{E}}^{\mathcal{E}'}$$

which corresponds to Theorem 11.3.

Remark that if  $U = V$  and  $\mathcal{E} = \mathcal{F}$ ,  $\mathcal{E}' = \mathcal{F}'$ , then the statement of Theorem 11.3 simply gives us that

$$[T]_{\mathcal{E}'}^{\mathcal{E}'} = [I_U]_{\mathcal{E}'}^{\mathcal{E}} [T]_{\mathcal{E}}^{\mathcal{E}} ([I_U]_{\mathcal{E}}^{\mathcal{E}'})^{-1}.$$

In other words,  $[T]_{\mathcal{E}'}^{\mathcal{E}'}$  is the conjugate of  $[T]_{\mathcal{E}}^{\mathcal{E}}$  by the change of bases matrix  $[I_U]_{\mathcal{E}}^{\mathcal{E}'}$ .

## 12 Similar Matrices, Eigenvectors and Eigenvalues

### 12.1 Similar matrices

In Section 11 we studied what happens to the matrix of a linear map  $T : U \rightarrow V$  when we change bases of  $U$  and  $V$ . Now we look at the case when  $U = V$ , where we only have a single vector space  $V$ , and a single change of basis. Surprisingly, this turns out to be more complicated than the situation with two different spaces.

Let  $V$  be a vector space of dimension  $n$  over the field  $K$ , and let  $T : V \rightarrow V$  be a linear map. Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  and  $\mathbf{e}'_1, \dots, \mathbf{e}'_n$  be two bases of  $V$ , and let  $A = (\alpha_{ij})$  and  $B = (\beta_{ij})$  be the matrices of  $T$  with respect to  $\{\mathbf{e}_i\}$  and  $\{\mathbf{e}'_i\}$  respectively. Let  $P = (\sigma_{ij})$  be the change of basis matrix from  $\{\mathbf{e}'_i\}$  to  $\{\mathbf{e}_i\}$ . *Note that this is the opposite change of basis to the one considered in the last section.*

Then applying Theorem 11.3 applies, but with both  $Q$  and  $P$  replaced by  $P^{-1}$  we have:

**Theorem 12.1** *With the notation above,  $B = P^{-1}AP$ .*

**Definition** Two  $n \times n$  matrices over  $K$  are said to be *similar* if there exists an  $n \times n$  invertible matrix  $P$  with  $B = P^{-1}AP$ .

So two matrices are similar if and only if they represent the same linear map  $T : V \rightarrow V$  with respect to different bases of  $V$ . It is easily checked that similarity is an equivalence relation on the set of  $n \times n$  matrices over  $K$ .

We saw in Theorem 11.5 that two matrices of the same size are equivalent if and only if they have the same rank. It is more difficult to decide whether two matrices are similar. Similar matrices are certainly equivalent, so they have the same rank, but equivalent matrices need not be similar.

**Example** Let  $A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ .

Then  $A$  and  $B$  both have rank 2, so they are equivalent. However, since  $A = I_2$ , for any invertible  $2 \times 2$  matrix  $P$  we have  $P^{-1}AP = A$ , so  $A$  is similar only to itself. Hence  $A$  and  $B$  are not similar.

To decide whether matrices are similar, it would be helpful to have a canonical form, just like we had the canonical form  $E_s$  in Section 11 for equivalence. Then we could test for similarity by reducing  $A$  and  $B$  to canonical form and checking whether we get the same result. But this turns out to be quite difficult, and depends on the field  $K$ . For the case  $K = \mathbb{C}$  (the complex numbers), we have the *Jordan Canonical Form*, which Maths students learn about in the Second Year.

## 12.2 Eigenvectors and eigenvalues

In this course, we shall only consider the question of which matrices are similar to a diagonal matrix. Such matrices are said to be *diagonalisable*. (Recall that  $A = (\alpha_{ij})$  is diagonal if  $\alpha_{ij} = 0$  for  $i \neq j$ .) We shall see, for example, that the matrix  $B$  in the example above is not diagonalisable.

It turns out that the possible entries on the diagonal of a matrix similar to  $A$  can be calculated directly from  $A$ . They are called *eigenvalues* of  $A$  and depend only on the linear map to which  $A$  corresponds, and not on the particular choice of basis.

**Definition** Let  $T : V \rightarrow V$  be a linear map, where  $V$  is a vector space over  $K$ . Suppose that for some non-zero vector  $\mathbf{v} \in V$  and some scalar  $\lambda \in K$ , we have  $T(\mathbf{v}) = \lambda\mathbf{v}$ . Then  $\mathbf{v}$  is called an *eigenvector* of  $T$ , and  $\lambda$  is called the *eigenvalue* of  $T$  corresponding to  $\mathbf{v}$ .

Note that the zero vector is **not** an eigenvector. (This would not be a good idea, because  $T\mathbf{0} = \lambda\mathbf{0}$  for all  $\lambda$ .) However, the zero scalar  $0_K$  may sometimes be an eigenvalue.

**Example** Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be defined by  $T(\alpha_1, \alpha_2) = (2\alpha_1, 0)$ . Then  $T(1, 0) = 2(1, 0)$ , so 2 is an eigenvalue and  $(1, 0)$  an eigenvector. Also  $T(0, 1) = (0, 0) = 0(0, 1)$ , so 0 is an eigenvalue and  $(0, 1)$  an eigenvector.

In this example, notice that in fact  $(\alpha, 0)$  and  $(0, \alpha)$  are eigenvectors for any  $\alpha \neq 0$ . In general, it is easy to see that if  $\mathbf{v}$  is an eigenvector of  $T$ , then so is  $\alpha\mathbf{v}$  for any non-zero scalar  $\alpha$ .

In some books, eigenvectors and eigenvalues are called *characteristic vectors* and *characteristic roots*, respectively.

Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  be a basis of  $V$ , and let  $A = (\alpha_{ij})$  be the matrix of  $T$  with respect to this basis. As in Section 7.1, to each vector  $\mathbf{v} = \lambda_1\mathbf{e}_1 + \dots + \lambda_n\mathbf{e}_n \in V$ , we associate the column vector

$$\underline{\mathbf{v}} = \begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \cdot \\ \cdot \\ \lambda_n \end{pmatrix} \in K^{n,1}.$$

Then, by Proposition 7.2, for  $\mathbf{u}, \mathbf{v} \in V$ , we have  $T(\mathbf{u}) = \mathbf{v}$  if and only if  $A\underline{\mathbf{u}} = \underline{\mathbf{v}}$ , and in particular

$$T(\mathbf{v}) = \lambda\mathbf{v} \iff A\underline{\mathbf{v}} = \lambda\underline{\mathbf{v}}.$$

**Definition** Let  $A$  be an  $n \times n$  matrix over  $K$ . Suppose that, for some non-zero column vector  $\underline{\mathbf{v}} \in K^{n,1}$  and some scalar  $\lambda \in K$ , we have  $A\underline{\mathbf{v}} = \lambda\underline{\mathbf{v}}$ . Then  $\underline{\mathbf{v}}$  is called an *eigenvector* of  $A$ , and  $\lambda$  is called the *eigenvalue* of  $A$  corresponding to  $\underline{\mathbf{v}}$ .

It follows from Proposition 7.2 that if the matrix  $A$  corresponds to the linear map  $T$ , then  $\lambda$  is an eigenvalue of  $T$  if and only if it is an eigenvalue of  $A$ . It follows immediately that similar matrices have the same eigenvalues, because they represent the same linear map with respect to different bases. We shall give another proof of this fact in Theorem 12.3 below.

**Theorem 12.2** Let  $A$  be an  $n \times n$  matrix. Then  $\lambda$  is an eigenvalue of  $A$  if and only if  $\det(A - \lambda I_n) = 0$ .

PROOF: Suppose that  $\lambda$  is an eigenvalue of  $A$ . Then  $A\underline{\mathbf{v}} = \lambda\underline{\mathbf{v}}$  for some non-zero  $\underline{\mathbf{v}} \in K^{n,1}$ . This is equivalent to  $A\underline{\mathbf{v}} = \lambda I_n \underline{\mathbf{v}}$ , or  $(A - \lambda I_n)\underline{\mathbf{v}} = \underline{\mathbf{0}}$ . But this says exactly that  $\underline{\mathbf{v}}$  is a non-zero solution to the homogeneous system of simultaneous equations defined by the matrix  $A - \lambda I_n$ , and then by Theorem 9.6(i),  $A - \lambda I_n$  is singular, and so  $\det(A - \lambda I_n) = 0$  by Theorem 10.5.

Conversely, if  $\det(A - \lambda I_n) = 0$  then  $A - \lambda I_n$  is singular, and so by Theorem 9.6(i) the system of simultaneous equations defined by  $A - \lambda I_n$  has nonzero solutions. Hence there exists a non-zero  $\underline{\mathbf{v}} \in K^{n,1}$  with  $(A - \lambda I_n)\underline{\mathbf{v}} = \underline{\mathbf{0}}$ , which is equivalent to  $A\underline{\mathbf{v}} = \lambda I_n \underline{\mathbf{v}}$ , and so  $\lambda$  is an eigenvalue of  $A$ .  $\square$

**Definition** For an  $n \times n$  matrix  $A$ , the equation  $\det(A - xI_n) = 0$  is called the *characteristic equation* of  $A$ , and  $\det(A - xI_n)$  is called the *characteristic polynomial* of  $A$ . It is a polynomial of degree  $n$  in  $x$ .

The above theorem says that the eigenvalues of  $A$  are the roots of the characteristic equation, which means that we have a method of calculating them. Once the eigenvalues are known, it is then straightforward to compute the corresponding eigenvectors.

**Example 1** Let  $A = \begin{pmatrix} 1 & 2 \\ 5 & 4 \end{pmatrix}$ . Then

$$\det(A - xI_2) = \begin{vmatrix} 1-x & 2 \\ 5 & 4-x \end{vmatrix} = (1-x)(4-x) - 10 = x^2 - 5x - 6 = (x-6)(x+1).$$

Hence the eigenvalues of  $A$  are the roots of  $(x-6)(x+1) = 0$ ; that is 6 and  $-1$ .

Let us now find the eigenvectors corresponding to the eigenvalue 6. We seek a non-zero column vector  $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$  such that

$$\begin{pmatrix} 1 & 2 \\ 5 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 6 \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}; \quad \text{that is} \quad \begin{pmatrix} -5 & 2 \\ 5 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

We can take  $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 2 \\ 5 \end{pmatrix}$  to be our eigenvector; or indeed any non-zero multiple of  $\begin{pmatrix} 2 \\ 5 \end{pmatrix}$ .

Similarly, for the eigenvalue  $-1$ , we want a non-zero column vector  $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$  such that

$$\begin{pmatrix} 1 & 2 \\ 5 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = -1 \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}; \quad \text{that is} \quad \begin{pmatrix} 2 & 2 \\ 5 & 5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix},$$

and we can take  $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$  to be our eigenvector.

**Example 2** This example shows that the eigenvalues can depend on the field  $K$ . Let

$$A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}. \quad \text{Then} \quad \det(A - xI_2) = \begin{vmatrix} -x & -1 \\ 1 & -x \end{vmatrix} = x^2 + 1,$$

so the characteristic equation is  $x^2 + 1 = 0$ . If  $K = \mathbb{R}$  (the real numbers) then this equation has no solutions, so there are no eigenvalues or eigenvectors. However, if  $K = \mathbb{C}$  (the complex numbers), then there are two eigenvalues  $i$  and  $-i$ , and by a similar calculation to the one in the last example, we find that  $\begin{pmatrix} -1 \\ i \end{pmatrix}$  and  $\begin{pmatrix} 1 \\ i \end{pmatrix}$  are eigenvectors corresponding to  $i$  and  $-i$  respectively.

**Theorem 12.3** *Similar matrices have the same characteristic equation and hence the same eigenvalues.*

PROOF: Let  $A$  and  $B$  be similar matrices. Then there exists an invertible matrix  $P$  with  $B = P^{-1}AP$ . Then

$$\det(B - xI_n) = \det(P^{-1}AP - xI_n) = \det(P^{-1}(A - xI_n)P) = (\text{by Theorem 10.3})$$

$$\det(P^{-1}) \det(A - xI_n) \det(P) = \det(P^{-1}) \det(P) \det(A - xI_n) = \det(A - xI_n).$$

Hence  $A$  and  $B$  have the same characteristic equation. Since the eigenvalues are the roots of the characteristic equation, they have the same eigenvalues.  $\square$

Since the different matrices corresponding to a linear map  $T$  are all equivalent, they all have the same characteristic equation, so we can unambiguously refer to it also as the characteristic equation of  $T$  if we want to.

There is one case where the eigenvalues can be written down immediately.

**Proposition 12.4** *Suppose that the matrix  $A$  is upper triangular. Then the eigenvalues of  $A$  are just the diagonal entries  $\alpha_{ii}$  of  $A$ .*

PROOF: We saw in Corollary 10.2 that the determinant of  $A$  is the product of the diagonal entries  $\alpha_{ii}$ . Hence the characteristic polynomial of such a matrix is  $\prod_{i=1}^n (\alpha_{ii} - x)$ , and so the eigenvalues are the  $\alpha_{ii}$ .  $\square$

**Example** Let  $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ . Then  $A$  is upper triangular, so its only eigenvalue is 1. We can now see that  $A$  cannot be similar to any diagonal matrix  $B$ . Such a  $B$  would also have just 1 as an eigenvalue, and then, by Proposition 10.2 again, this would force  $B$  to be the identity matrix  $I_2$ . But  $P^{-1}I_2P = I_2$  for any invertible matrix  $P$ , so  $I_2$  is not similar to any matrix other than itself! So  $A$  cannot be similar to  $I_2$ , and hence  $A$  is not diagonalisable.

The next theorem describes the connection between diagonalisable matrices and eigenvectors. If you have understood everything so far then its proof should be almost obvious.

**Theorem 12.5** *Let  $T : V \rightarrow V$  be a linear map. Then the matrix of  $T$  is diagonal with respect to some basis of  $V$  if and only if  $V$  has a basis consisting of eigenvectors of  $T$ .*

*Equivalently, let  $A$  be an  $n \times n$  matrix over  $K$ . Then  $A$  is similar to a diagonal matrix if and only if the space  $K^{n,1}$  has a basis of eigenvectors of  $A$ .*

PROOF: The equivalence of the two statements follows directly from the correspondence between linear maps and matrices, and the corresponding definitions of eigenvectors and eigenvalues.

Suppose that the matrix  $A = (\alpha_{ij})$  of  $T$  is diagonal with respect to the basis  $\mathbf{e}_1, \dots, \mathbf{e}_n$  of  $V$ . Recall from Section 7.1 that the images of the  $i$ -th basis vector of  $V$  is represented by the  $i$ -th column of  $A$ . But since  $A$  is diagonal, this column has the single non-zero entry  $\alpha_{ii}$ . Hence  $T(\mathbf{e}_i) = \alpha_{ii}\mathbf{e}_i$ , and so each basis vector  $\mathbf{e}_i$  is an eigenvector of  $A$ .

Conversely, suppose that  $\mathbf{e}_1, \dots, \mathbf{e}_n$  is a basis of  $V$  consisting entirely of eigenvectors of  $T$ . Then, for each  $i$ , we have  $T(\mathbf{e}_i) = \lambda_i \mathbf{e}_i$  for some  $\lambda_i \in K$ . But then the matrix of  $A$  with respect to this basis is the diagonal matrix  $A = (\alpha_{ij})$  with  $\alpha_{ii} = \lambda_i$  for each  $i$ .  $\square$

We now show that  $A$  is diagonalisable in the case when there are  $n$  distinct eigenvalues.

**Theorem 12.6** *Let  $\lambda_1, \dots, \lambda_r$  be distinct eigenvalues of  $T : V \rightarrow V$ , and let  $\mathbf{v}_1, \dots, \mathbf{v}_r$  be corresponding eigenvectors. (So  $T(\mathbf{v}_i) = \lambda_i \mathbf{v}_i$  for  $1 \leq i \leq r$ .) Then  $\mathbf{v}_1, \dots, \mathbf{v}_r$  are linearly independent.*

PROOF: We prove this by induction on  $r$ . It is true for  $r = 1$ , because eigenvectors are non-zero by definition. For  $r > 1$ , suppose that for some  $\alpha_1, \dots, \alpha_r \in K$  we have

$$\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_r \mathbf{v}_r = \mathbf{0}.$$

Then, applying  $T$  to this equation gives

$$\alpha_1 \lambda_1 \mathbf{v}_1 + \alpha_2 \lambda_2 \mathbf{v}_2 + \dots + \alpha_r \lambda_r \mathbf{v}_r = \mathbf{0}.$$

Now, subtracting  $\lambda_1$  times the first equation from the second gives

$$\alpha_2 (\lambda_2 - \lambda_1) \mathbf{v}_2 + \dots + \alpha_r (\lambda_r - \lambda_1) \mathbf{v}_r = \mathbf{0}.$$

By inductive hypothesis,  $\mathbf{v}_2, \dots, \mathbf{v}_r$  are linearly independent, so  $\alpha_i (\lambda_i - \lambda_1) = 0$  for  $2 \leq i \leq r$ . But, by assumption,  $\lambda_i - \lambda_1 \neq 0$  for  $i > 1$ , so we must have  $\alpha_i = 0$  for  $i > 1$ . But then  $\alpha_1 \mathbf{v}_1 = \mathbf{0}$ , so  $\alpha_1$  is also zero. Thus  $\alpha_i = 0$  for all  $i$ , which proves that  $\mathbf{v}_1, \dots, \mathbf{v}_r$  are linearly independent.  $\square$

**Corollary 12.7** *If the linear map  $T : V \rightarrow V$  (or equivalently the  $n \times n$  matrix  $A$ ) has  $n$  distinct eigenvalues, where  $n = \dim(V)$ , then  $T$  (or  $A$ ) is diagonalisable.*

PROOF: Under the hypothesis, there are  $n$  linearly independent eigenvectors, which form a basis of  $V$  by Corollary 3.9. The result follows from Theorem 12.5.  $\square$

### Example

$$A = \begin{pmatrix} 4 & 5 & 2 \\ -6 & -9 & -4 \\ 6 & 9 & 4 \end{pmatrix}. \quad \text{Then } |A - xI_3| = \begin{vmatrix} 4-x & 5 & 2 \\ -6 & -9-x & -4 \\ 6 & 9 & 4-x \end{vmatrix}.$$

To help evaluate this determinant, apply first the row operation  $\mathbf{r}_3 \rightarrow \mathbf{r}_3 + \mathbf{r}_2$  and then the column operation  $\mathbf{c}_2 \rightarrow \mathbf{c}_2 - \mathbf{c}_3$ , giving

$$|A - xI_3| = \begin{vmatrix} 4-x & 5 & 2 \\ -6 & -9-x & -4 \\ 0 & -x & -x \end{vmatrix} = \begin{vmatrix} 4-x & 3 & 2 \\ -6 & -5-x & -4 \\ 0 & 0 & -x \end{vmatrix},$$

and then expanding by the third row we get

$$-x((4-x)(-5-x) + 18) = -x(x^2 + x - 2) = -x(x+2)(x-1)$$

so the eigenvalues are 0, 1 and  $-2$ . Since these are distinct, we know from the above corollary that  $A$  can be diagonalised. In fact, the eigenvectors will be the new basis for the diagonal matrix, so we will calculate these.

In the following calculations, we will denote eigenvectors  $\underline{\mathbf{v}}_1$ , etc. by  $\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$ , where  $x_1, x_2, x_3$  need to be calculated by solving simultaneous equations.

For the eigenvalue  $\lambda = 0$ , an eigenvector  $\underline{\mathbf{v}}_1$  satisfies  $A\underline{\mathbf{v}}_1 = \underline{\mathbf{0}}$ , which gives the three equations:

$$4x_1 + 5x_2 + 2x_3 = 0; \quad -6x_1 - 9x_2 - 4x_3 = 0; \quad 6x_1 + 9x_2 + 4x_3 = 0.$$

The third is clearly redundant, and adding twice the first to the second gives  $2x_1 + x_2 = 0$

and then we see that one solution is  $\underline{\mathbf{v}}_1 = \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix}$ .

For  $\lambda = 1$ , we want an eigenvector  $\underline{\mathbf{v}}_2$  with  $A\underline{\mathbf{v}}_2 = \underline{\mathbf{v}}_2$ , which gives the equations

$$\begin{aligned} 4x_1 + 5x_2 + 2x_3 = x_1; & \quad -6x_1 - 9x_2 - 4x_3 = x_2; & \quad 6x_1 + 9x_2 + 4x_3 = x_3; & \quad \implies \\ 3x_1 + 5x_2 + 2x_3 = 0; & \quad -6x_1 - 10x_2 - 4x_3 = 0; & \quad 6x_1 + 9x_2 + 3x_3 = 0. \end{aligned}$$

Adding the second and third equations gives  $x_2 + x_3 = 0$  and then we see that a solution is

$$\underline{\mathbf{v}}_2 = \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}.$$

Finally, for  $\lambda = -2$ ,  $A\underline{\mathbf{v}}_3 = -2\underline{\mathbf{v}}_3$  gives the equations

$$6x_1 + 5x_2 + 2x_3 = 0; \quad -6x_1 - 7x_2 - 4x_3 = 0; \quad 6x_1 + 9x_2 + 6x_3 = 0,$$

of which one solution is  $\underline{\mathbf{v}}_3 = \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}$ .

Now, if we change basis to  $\underline{\mathbf{v}}_1, \underline{\mathbf{v}}_2, \underline{\mathbf{v}}_3$ , we should get the diagonal matrix with the eigenvalues  $0, 1, -2$  on the diagonal. We can check this by direct calculation. Remember that  $P$  is the change of basis matrix from the new basis to the old one and has columns the new basis vectors expressed in terms of the old. But the old basis is the standard basis, so the columns of  $P$  are the new basis vectors. Hence

$$P = \begin{pmatrix} 1 & 1 & 1 \\ -2 & -1 & -2 \\ 3 & 1 & 2 \end{pmatrix} \text{ and, according to Theorem 12.1, we should have } P^{-1}AP = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}.$$

To check this, we first need to calculate  $P^{-1}$ , either by row reduction or by the cofactor method. The answer turns out to be  $P^{-1} = \begin{pmatrix} 0 & 1 & 1 \\ 2 & 1 & 0 \\ -1 & -2 & -1 \end{pmatrix}$ , and now we can check that

the above equation really does hold.

**Warning** The converse of Corollary 12.7 is not true. If it turns out that there do not exist  $n$  distinct eigenvalues, then you cannot conclude from this that the matrix is not be diagonalisable. This is really rather obvious, because the identity matrix has only a single eigenvalue, but it is diagonal already. Even so, this is one of the most common mistakes that students make.

If there are fewer than  $n$  distinct eigenvalues, then the matrix may or may not be diagonalisable, and you have to test directly to see whether there are  $n$  linearly independent eigenvectors. Let us consider two rather similar looking examples.

$$A_1 = \begin{pmatrix} 1 & 1 & 1 \\ 0 & -1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 1 & 2 & -2 \\ 0 & -1 & 2 \\ 0 & 0 & 1 \end{pmatrix}.$$

Both matrices are upper triangular, so we know from Proposition 12.4 that both have eigenvalues  $1$  and  $-1$ , with  $1$  repeated. Since  $-1$  occurs only once, it can only have a single

associated linearly independent eigenvector. (Can you prove that?) Solving the equations as usual, we find that  $A_1$  and  $A_2$  have eigenvectors  $\begin{pmatrix} 1 \\ -2 \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$ , respectively, associated with eigenvalue  $-1$ .

The repeated eigenvalue 1 is more interesting, because there could be one or two associated linearly independent eigenvectors. The equation  $A_1\mathbf{x} = \mathbf{x}$  gives the equations

$$x_1 + x_2 + x_3 = x_1; \quad -x_2 + x_3 = x_2; \quad x_3 = x_3,$$

so  $x_2 + x_3 = -2x_2 + x_3 = 0$ , which implies that  $x_2 = x_3 = 0$ . Hence the only eigenvectors are multiples of  $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ . Hence  $A_1$  has only two linearly independent eigenvectors in total, and so it cannot be diagonalised.

On the other hand,  $A_2\mathbf{x} = \mathbf{x}$  gives the equations

$$x_1 + 2x_2 - 2x_3 = x_1; \quad -x_2 + 2x_3 = x_2; \quad x_3 = x_3,$$

which reduce to the single equation  $x_2 - x_3 = 0$ . This time there are two linearly independent solutions, giving eigenvectors  $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$ . So  $A_2$  has three linearly independent eigenvectors in total, and it can be diagonalised. In fact, using the eigenvectors as columns of the change of basis matrix  $P$  as before gives

$$P = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \text{ and then we find that } P^{-1} = \begin{pmatrix} 1 & 1 & -1 \\ 0 & 0 & 1 \\ 0 & -1 & 1 \end{pmatrix}, \text{ and we can check that}$$

$$P^{-1}A_2P = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

### 12.3 The scalar product - symmetric and orthogonal matrices

**Definition** The (standard) *scalar product* of two vectors  $\mathbf{v} = (\alpha_1, \dots, \alpha_n)$  and  $\mathbf{w} = (\beta_1, \dots, \beta_n)$  in  $\mathbb{R}^n$  is defined to be

$$\mathbf{v} \cdot \mathbf{w} = \sum_{i=1}^n \alpha_i \beta_i.$$

**Definition** A basis  $\mathbf{b}_1, \dots, \mathbf{b}_n$  of  $\mathbb{R}^n$  is called *orthonormal* if

- (i)  $\mathbf{b}_i \cdot \mathbf{b}_i = 1$  for  $1 \leq i \leq n$ , and
- (ii)  $\mathbf{b}_i \cdot \mathbf{b}_j = 0$  for  $1 \leq i, j \leq n$  and  $i \neq j$ .

In other words, an orthonormal basis consists of mutually orthogonal vectors of length 1. For example, the standard basis is orthonormal.

**Definition** An  $n \times n$  matrix  $A$  is said to *symmetric* if  $A^T = A$ .

**Definition** An  $n \times n$  matrix  $A$  is said to *orthogonal* if  $A^T = A^{-1}$  or, equivalently, if  $AA^T = A^T A = I_n$ .

**Examples.**

$$\begin{pmatrix} 2/\sqrt{13} & -3/\sqrt{13} \\ 3/\sqrt{13} & 2/\sqrt{13} \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1/3 & 2/3 & 2/3 \\ 2/3 & -2/3 & 1/3 \\ 2/3 & 1/3 & -2/3 \end{pmatrix}$$

are both orthogonal matrices.

The main result of this section is that we can diagonalise any real symmetric matrix  $A$  by a real orthogonal matrix. We shall prove this only in the case when  $A$  has distinct eigenvalues; the complete proof will be given in Year 2.

**Proposition 12.8** *An  $n \times n$  matrix  $A$  over  $\mathbb{R}$  is orthogonal if and only if the rows  $\mathbf{r}_1, \dots, \mathbf{r}_n$  of  $A$  form an orthonormal basis of  $\mathbb{R}^n$  if and only if the columns  $\mathbf{c}_1, \dots, \mathbf{c}_n$  of  $A$  form an orthonormal basis of  $\mathbb{R}^{n,1}$ .*

PROOF: Note that an orthogonal matrix  $A$  is invertible, which by Theorem 9.2 implies that its row and column ranks are equal to  $n$ , and hence that the rows of  $A$  form a basis of  $\mathbb{R}^n$  and the columns form a basis of  $\mathbb{R}^{n,1}$ . By the definition of matrix multiplication,  $AA^T = I_n$  implies that  $\mathbf{r}_i \cdot \mathbf{r}_i = 1$  and  $\mathbf{r}_i \cdot \mathbf{r}_j = 0$  for  $i \neq j$ , and hence that the rows form an orthonormal basis of  $\mathbb{R}^n$ . Similarly,  $A^T A = I_n$  implies that the columns of  $A$  form an orthonormal basis of  $\mathbb{R}^{n,1}$ . Conversely, if the rows or columns of  $A$  form an orthonormal basis of  $\mathbb{R}^n$  or  $\mathbb{R}^{n,1}$ , then we get  $AA^T = I_n$  or  $A^T A = I_n$ , both of which imply that  $A^T = A^{-1}$ ; that is, that  $A$  is orthogonal.  $\square$

**Proposition 12.9** *Let  $A$  be a real symmetric matrix. Then  $A$  has an eigenvalue in  $\mathbb{R}$ , and all complex eigenvalues of  $A$  lie in  $\mathbb{R}$ .*

PROOF: (To simplify the notation, we will write just  $\mathbf{v}$  for a column vector  $\underline{\mathbf{v}}$  in this proof.)

The characteristic equation  $\det(A - xI_n) = 0$  is a polynomial equation of degree  $n$  in  $x$ , and since  $\mathbb{C}$  is an algebraically closed field, it certainly has a root  $\lambda \in \mathbb{C}$ , which is an eigenvalue for  $A$  if we regard  $A$  as a matrix over  $\mathbb{C}$ . We shall prove that any such  $\lambda$  lies in  $\mathbb{R}$ , which will prove the proposition.

For a column vector  $\mathbf{v}$  or matrix  $B$  over  $\mathbb{C}$ , we denote by  $\bar{\mathbf{v}}$  or  $\bar{B}$  the result of replacing all entries of  $\mathbf{v}$  or  $B$  by their complex conjugates. Since the entries of  $A$  lie in  $\mathbb{R}$ , we have  $\bar{A} = A$ .

Let  $\mathbf{v}$  be a complex eigenvector associated with  $\lambda$ . Then

$$A\mathbf{v} = \lambda\mathbf{v} \quad (1)$$

so, taking complex conjugates and using  $\bar{A} = A$ , we get

$$A\bar{\mathbf{v}} = \bar{\lambda}\bar{\mathbf{v}}. \quad (2)$$

Transposing (1) and using  $A^T = A$  gives

$$\mathbf{v}^T A = \lambda\mathbf{v}^T, \quad (3)$$

so by (2) and (3) we have

$$\lambda\mathbf{v}^T \bar{\mathbf{v}} = \mathbf{v}^T A \bar{\mathbf{v}} = \bar{\lambda}\mathbf{v}^T \bar{\mathbf{v}}.$$

But if  $\mathbf{v} = (\alpha_1, \alpha_2, \dots, \alpha_n)^T$ , then  $\mathbf{v}^T \bar{\mathbf{v}} = \alpha_1 \bar{\alpha}_1 + \dots + \alpha_n \bar{\alpha}_n$ , which is a nonzero real number (eigenvectors are nonzero by definition). Thus  $\lambda = \bar{\lambda}$ , so  $\lambda \in \mathbb{R}$ .  $\square$

**Proposition 12.10** *Let  $A$  be a real symmetric matrix, and let  $\lambda_1, \lambda_2$  be two distinct eigenvalues of  $A$ , with corresponding eigenvectors  $\mathbf{v}_1, \mathbf{v}_2$ . Then  $\mathbf{v}_1 \cdot \mathbf{v}_2 = 0$ .*

PROOF: (As in Proposition 12.9, we will write  $\mathbf{v}$  rather than  $\underline{\mathbf{v}}$  for a column vector in this proof. So  $\mathbf{v}_1 \cdot \mathbf{v}_2$  is the same as  $\mathbf{v}_1^T \mathbf{v}_2$ .) We have

$$A\mathbf{v}_1 = \lambda_1\mathbf{v}_1 \quad (1) \quad \text{and} \quad A\mathbf{v}_2 = \lambda_2\mathbf{v}_2 \quad (2).$$

Transposing (1) and using  $A = A^T$  gives  $\mathbf{v}_1^T A = \lambda_1 \mathbf{v}_1^T$ , and so

$$\mathbf{v}_1^T A \mathbf{v}_2 = \lambda_1 \mathbf{v}_1^T \mathbf{v}_2 \quad (3) \quad \text{and similarly} \quad \mathbf{v}_2^T A \mathbf{v}_1 = \lambda_2 \mathbf{v}_2^T \mathbf{v}_1 \quad (4).$$

Transposing (4) gives  $\mathbf{v}_1^T A \mathbf{v}_2 = \lambda_2 \mathbf{v}_1^T \mathbf{v}_2$  and subtracting (3) from this gives  $(\lambda_2 - \lambda_1) \mathbf{v}_1^T \mathbf{v}_2 = 0$ . Since  $\lambda_2 - \lambda_1 \neq 0$  by assumption, we have  $\mathbf{v}_1^T \mathbf{v}_2 = 0$ .  $\square$

**Theorem 12.11** *Let  $A$  be a real symmetric  $n \times n$  matrix. Then there exists a real orthogonal matrix  $P$  with  $P^{-1}AP (= P^TAP)$  diagonal.*

PROOF: We shall prove this only in the case when the eigenvalues  $\lambda_1, \dots, \lambda_n$  of  $A$  are all distinct. By Proposition 12.9 we have  $\lambda_i \in \mathbb{R}$  for all  $i$ , and so there exist associated eigenvectors  $\mathbf{v}_i \in \mathbb{R}^{n,1}$ . By Proposition 12.10, we have  $\mathbf{v}_i \cdot \mathbf{v}_j = \mathbf{v}_i^T \mathbf{v}_j = 0$  for  $i \neq j$ . Since each  $\mathbf{v}_i$  is non-zero, we have  $\mathbf{v}_i \cdot \mathbf{v}_i = \alpha_i > 0$ . By replacing each  $\mathbf{v}_i$  by  $\mathbf{v}_i / \sqrt{\alpha_i}$  (which is also an eigenvector for  $\lambda_i$ ), we can assume that  $\mathbf{v}_i \cdot \mathbf{v}_i = 1$  for all  $i$ . Since, by Theorem 12.6, the  $\mathbf{v}_i$  are linearly independent, they form a basis and hence an orthonormal basis of  $\mathbb{R}^{n,1}$ . So, by Proposition 12.8, the matrix  $P$  with columns  $\mathbf{v}_1, \dots, \mathbf{v}_n$  is orthogonal. But  $P^{-1}AP$  is the diagonal matrix with entries  $\lambda_1, \dots, \lambda_n$ , which proves the result.  $\square$

**Example.** Let

$$A = \begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix}.$$

Then

$$\det(A - \lambda I_2) = (1 - \lambda)^2 - 9 = \lambda^2 - 2\lambda - 8 = (\lambda - 4)(\lambda + 2),$$

so the eigenvalues of  $A$  are 4 and  $-2$ . Solving  $A\mathbf{v} = \lambda\mathbf{v}$  for  $\lambda = 4$  and  $-2$ , we find corresponding eigenvectors  $(1, 1)^T$  and  $(1, -1)^T$ . Proposition 12.10 tells us that these vectors are orthogonal to each other (which we can of course check directly!). Their lengths are both  $\sqrt{2}$ , so we divide by them by their lengths to give eigenvectors  $(1/\sqrt{2}, 1/\sqrt{2})^T$  and  $(1/\sqrt{2}, -1/\sqrt{2})^T$  of length 1.

The basis change matrix  $P$  has these vectors as columns, so  $P = \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix}$ , and we can check that  $P^T P = I_2$  (i.e.  $P$  is orthogonal) and that

$$P^{-1}AP = P^TAP = \begin{pmatrix} 4 & 0 \\ 0 & -2 \end{pmatrix}.$$

## 12.4 Linear recursive sequences

For our final topic we apply eigenvectors to find formulae for some linear recursive sequences. Let us look at a sequence  $z_i \in K$  defined by a linear recursion

$$z_0 = a_0, z_1 = a_1, \dots, z_{n-1} = a_{n-1}; \quad z_m = \sum_{i=0}^{n-1} \alpha_i z_{m-n+i}, \quad m \geq n \quad (4)$$

where the scalars  $\alpha_i, a_i \in K, i = 0, 1, \dots, n-1$  are given upfront. The most famous example of a linear recursive sequence is Fibonacci numbers

$$z_0 = 0, z_1 = 1; \quad z_m = z_{m-1} + z_{m-2}, \quad m \geq 2.$$

The question we would like to address is how to find an explicit (non-recursive) formula for  $z_m$ . Let us introduce a sequence of vectors in  $\mathbf{v}_m \in K^{n,1}$  and a matrix  $A \in K^{n,n}$ :

$$\mathbf{v}_m = \begin{pmatrix} z_{m+n-1} \\ z_{m+n-2} \\ \vdots \\ z_{m+2} \\ z_{m+1} \\ z_m \end{pmatrix}, \quad \mathbf{b} = \begin{pmatrix} a_{n-1} \\ a_{n-2} \\ \vdots \\ a_2 \\ a_1 \\ a_0 \end{pmatrix}, \quad A = \begin{pmatrix} \alpha_{n-1} & \alpha_{n-2} & \cdots & \alpha_2 & \alpha_1 & \alpha_0 \\ 1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 & 0 \\ \vdots & & & & & \vdots \\ 0 & 0 & \cdots & 1 & 0 & 0 \\ 0 & 0 & \cdots & 0 & 1 & 0 \end{pmatrix}. \quad (5)$$

If we can find an explicit formula for  $\mathbf{v}_m$  then we can find an explicit formula for  $z_m$ . On the other hand recursive relations (4) are equivalent to a single relation

$$\mathbf{v}_0 = \mathbf{b}, \quad \mathbf{v}_m = A\mathbf{v}_{m-1} \quad (6)$$

that results in an answer

$$\mathbf{v}_m = A^m \mathbf{b}. \quad (7)$$

We still have to compute  $\mathbf{v}_m$  explicitly in terms of  $\alpha_i$  and  $a_i$ . The following theorem is helpful. The first statement of the theorem is a very special feature of the matrices of linear recursive sequences. In fact, such a matrix  $A$  is already in a certain canonical form, called *Sylvester normal form*.

**Theorem 12.12** *The characteristic polynomial of  $A$  is  $(-1)^n(x^n - \sum_{k=0}^{n-1} \alpha_k x^k)$ . If  $\mathbf{b} = \mathbf{w}_1 + \mathbf{w}_2 + \dots + \mathbf{w}_s$  where each  $\mathbf{w}_i$  is an eigenvector with an eigenvalue  $\lambda_i$  then  $\mathbf{v}_m = \lambda_1^m \mathbf{w}_1 + \lambda_2^m \mathbf{w}_2 + \dots + \lambda_s^m \mathbf{w}_s$ .*

PROOF: The first statement is by induction. It is clear if  $n = 1$ . Having established the first statement for  $n - 1$ , we expand over the last column,

$$\begin{aligned} \det(A - xI_n) &= \begin{vmatrix} \alpha_{n-1} - x & \alpha_{n-2} & \cdots & \alpha_2 & \alpha_1 & \alpha_0 \\ 1 & -x & \cdots & 0 & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 & 0 \\ \vdots & & & & & \vdots \\ 0 & 0 & \cdots & 1 & -x & 0 \\ 0 & 0 & \cdots & 0 & 1 & -x \end{vmatrix} = \\ & -x \begin{vmatrix} \alpha_{n-1} - x & \alpha_{n-2} & \cdots & \alpha_2 & \alpha_1 \\ 1 & -x & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & & & & \vdots \\ 0 & 0 & \cdots & -x & 0 \\ 0 & 0 & \cdots & 1 & -x \end{vmatrix} + (-1)^{n+1} \alpha_0 \begin{vmatrix} 1 & -x & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & & & & \vdots \\ 0 & 0 & \cdots & 1 & -x \\ 0 & 0 & \cdots & 0 & 1 \end{vmatrix} \\ & = -x((-1)^{n-1}(x^{n-1} - \sum_{k=0}^{n-2} \alpha_{k+1} x^k)) + (-1)^{n+1} \alpha_0 = (-1)^n (x^n - \sum_{k=0}^{n-1} \alpha_k x^k). \end{aligned}$$

The second statement is clear,  $\mathbf{v}_m = A^m \mathbf{b} = A^m \mathbf{w}_1 + \dots + A^m \mathbf{w}_s = \lambda_1^m \mathbf{w}_1 + \lambda_2^m \mathbf{w}_2 + \dots + \lambda_s^m \mathbf{w}_s$ .  $\square$

For instance, for Fibonacci numbers

$$A = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \quad \mathbf{b} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

By Theorem 12.12, the characteristic polynomial of  $A$  is  $x^2 - x - 1$ , whose roots are  $\lambda_1 = (1 + \sqrt{5})/2$  and  $\lambda_2 = (1 - \sqrt{5})/2$ . Solving two linear systems  $(A - \lambda_i)\mathbf{x} = \mathbf{0}$  gives eigenvectors

$$\mathbf{w}_1 = \begin{pmatrix} 1 + \sqrt{5} \\ 2 \end{pmatrix}, \quad \mathbf{w}_2 = \begin{pmatrix} 1 - \sqrt{5} \\ 2 \end{pmatrix}.$$

Observe that  $\mathbf{b} = 1/2\sqrt{5}(\mathbf{w}_1 - \mathbf{w}_2)$ . Using Theorem 12.12, we conclude that

$$\mathbf{v}_m = \frac{1}{2\sqrt{5}} \left( \left( \frac{1 + \sqrt{5}}{2} \right)^m \mathbf{w}_1 - \left( \frac{1 - \sqrt{5}}{2} \right)^m \mathbf{w}_2 \right).$$

Looking at the lower entry, we arrive at Binet's formula,

$$z_m = \frac{1}{2\sqrt{5}} \left( \left( \frac{1 + \sqrt{5}}{2} \right)^m 2 - \left( \frac{1 - \sqrt{5}}{2} \right)^m 2 \right) = \frac{1}{\sqrt{5}} \left( \left( \frac{1 + \sqrt{5}}{2} \right)^m - \left( \frac{1 - \sqrt{5}}{2} \right)^m \right).$$