

# MA106 – Linear Algebra

## Assignment 1

January 2017

Answer the questions on your own paper. Write your own name in the top left-hand corner, and your supervisor's name in the top right-hand corner. Solutions to Problems 1,3,4,5 only must be handed in by **2.00 pm on Thursday 19 January** (Thursday of the second week of term), or they will not be marked.

**1.** Some of the following expressions define one or more numbers. Decide whether these lie in  $\mathbb{N}$ ,  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{R}$  or  $\mathbb{C}$ , or whether the expression does not define any numbers at all. (Of course, in some cases they will lie in more than one of these sets.)

$$0.00010001\dots \quad \text{where the '0001' repeats infinitely often,} \\ \sqrt{-4}, \quad \sqrt{6889}, \quad e^{6i\pi}, \quad 0^{-1}. \quad [5 \text{ marks}]$$

**2.** Many of the familiar properties of numbers can be deduced from the axioms A1–A4, M1–M4 and D. For example, the cancellation law of addition,  $\alpha + \beta = \alpha + \gamma \Rightarrow \beta = \gamma$  is proved by adding  $-\alpha$  to both sides of the equation and using A2; and the rule  $0\alpha = 0$  for all  $\alpha$  can be proved as follows:

$$0\alpha + 0 = 0\alpha \quad (\text{by A3}) = (0 + 0)\alpha \quad (\text{by A3}) = 0\alpha + 0\alpha \quad (\text{by D}),$$

and then  $0\alpha = 0$  follows from the cancellation law of addition.

Prove the following rules from the axioms (or from the two rules just proved above), for all numbers  $\alpha, \beta$ :

- (i)  $\alpha(-\beta) = (-\alpha)\beta = -(\alpha\beta)$ ;
- (ii)  $(-\alpha)(-\beta) = \alpha\beta$ ;
- (iii)  $(-1)\alpha = -\alpha$ .

**3.** Let  $\alpha$  and  $\beta$  be elements of a field, and suppose that  $\alpha\beta = 0$ . Prove that at least one of  $\alpha$  and  $\beta$  must equal 0. (You may assume the rule  $0\alpha = 0$  mentioned in Question 2.) [3 marks]

**4.** Let  $K$  be the set of complex numbers of the form  $a + ib\sqrt{17}$ , where  $a, b \in \mathbb{Q}$  (that is,  $a$  and  $b$  are rational numbers). Show that  $K$  is a “subfield” of  $\mathbb{C}$ , that is,  $1 \in K$  and  $\alpha - \beta, \alpha\beta, \gamma^{-1} \in K$  whenever  $\alpha, \beta \in K$  and  $\gamma \in K \setminus \{0\}$ . (It follows that  $K$  is itself a field but you don't need to prove this.) [7 marks]

**5.** Define a *paddock* to be a set in which A1–A4, M1–M4 and D hold, but instead of  $0 \neq 1$ , we have  $0 = 1$ . Find an example of a paddock. Prove that every paddock has just one element. Clearly the definition of paddock is of no use outside this exercise. [5 marks]

6. Let  $K$  be a field and  $\alpha_1, \dots, \alpha_n \in K$  where  $n \geq 1$ . Using the axioms of fields only, the sum

$$\alpha_1 + \alpha_2 + \cdots + \alpha_n \tag{1}$$

is meaningless, not because of the dots, but because only a sum of two elements is known to be defined! In this exercise we repair this.

Define  $\langle \alpha_1, \dots, \alpha_n \rangle$  recursively by  $\langle \alpha \rangle := \alpha$  and

$$\langle \alpha_1, \dots, \alpha_n \rangle := \langle \alpha_1, \dots, \alpha_{n-1} \rangle + \alpha_n \quad \text{if } n \geq 2.$$

(a) Prove

$$\langle \alpha_1, \dots, \alpha_k \rangle + \langle \beta_1, \dots, \beta_\ell \rangle = \langle \alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_\ell \rangle$$

for all  $\alpha_i, \beta_j \in K$  and all  $k, \ell \geq 1$ . Where are you using which axiom for fields? Hint: use induction.

Of course (1) is one of the more usual notations for  $\langle \alpha_1, \dots, \alpha_n \rangle$ .

(b) Let  $s$  be a bijective map from  $\{1, \dots, n\}$  to itself. Prove

$$\langle \alpha_1, \dots, \alpha_n \rangle = \langle \alpha_{s(1)}, \dots, \alpha_{s(n)} \rangle.$$

7. Let  $V$  be a vector space over a field  $K$ . Deduce the following facts from the vector space axioms. (The first one is done for you!)

(i)  $\alpha \mathbf{0} = \mathbf{0}$  for all  $\alpha \in K$ . Here  $\mathbf{0}$  is the zero vector;

(Solution:  $\alpha \mathbf{0} = \alpha(\mathbf{0} + \mathbf{0}) = \alpha \mathbf{0} + \alpha \mathbf{0}$ , so

$$\mathbf{0} = -(\alpha \mathbf{0}) + \alpha \mathbf{0} = -(\alpha \mathbf{0}) + \alpha \mathbf{0} + \alpha \mathbf{0} = \mathbf{0} + \alpha \mathbf{0} = \alpha \mathbf{0}.)$$

(ii)  $0 \mathbf{v} = \mathbf{0}$  for all  $\mathbf{v} \in V$ . Here  $0$  is the zero element of  $K$ ;

(iii)  $-(\alpha \mathbf{v}) = (-\alpha) \mathbf{v} = \alpha(-\mathbf{v})$ , for all  $\alpha \in K$  and  $\mathbf{v} \in V$ .

(iv) If  $\alpha \mathbf{v} = \mathbf{0}$  with  $\alpha \in K$ ,  $\mathbf{v} \in V$ , then either  $\alpha = 0$  or  $\mathbf{v} = \mathbf{0}$ .